

# Mysterious pulsar with hidden powers discovered

October 14 2010

---




Illustration: NASA/CXC/M.Weiss

Dramatic flares and bursts of energy - activity previously thought reserved for only the strongest magnetized pulsars - has been observed emanating from a weakly magnetised, slowly rotating pulsar. The international team of astrophysicists who made the discovery believe that the source of the pulsar's power may be hidden deep within its surface.

Pulsars, or [neutron stars](#), are the collapsed remains of [massive stars](#). Although they are on average only about 30km in diameter, they have hugely powerful surface magnetic fields, billions of times that of our Sun.

The most extreme kind of pulsars have a surface [magnetic field](#) 50-1000 times stronger than normal and emit powerful flares of gamma rays and X-rays. Named magnetars (which stands for "magnetic stars") by astronomers, their huge magnetic fields are thought to be the ultimate source of power for the bursts of [gamma rays](#).

Theoretical studies indicate that in magnetars the internal field is actually stronger than the surface field, a property which can deform the crust and propagate outwards. The decay of the magnetic field leads to the production of steady and bursting X-ray emission through the heating of the neutron star crust or the acceleration of particles.

Now, research published today in [Science Express](#), suggests that the same power source can also work for weaker, non-magnetar, pulsars. The observations, which were made by NASA's Chandra and Swift X-ray observatories of the neutron star SGR 0418, may indicate the presence of a huge internal magnetic field in these seemingly less powerful pulsars, which is not matched by their surface magnetic field.

"We have now discovered bursts and flares, i.e. magnetar-like activity, from a new [pulsar](#) whose magnetic field is very low," said Dr Silvia Zane, from UCL's (University College London) Mullard Space Science Laboratory, and an author of the research.

Pulsars are highly magnetized, and as they rotate winds of high-energy particles carry energy away from the star, causing the rotation rate of the star to gradually decrease. What sets SGR 0418 apart from similar neutron stars is that, unlike those stars that are observed to be gradually rotating more slowly, careful monitoring of SGR 0418 over a span of 490 days has revealed no evidence that its rotation is decreasing.

"It is the very first time this has been observed and the discovery poses the question of where the powering mechanism is in this case. At this

point, we are also interested in how many of the other normal, low field neutron stars that populate the galaxy can at some point wake up and manifest themselves as a flaring source," said Dr Zane.

A crucial question is how large an imbalance can be maintained between the surface and interior magnetic fields. SGR 0418 represents an important test case.

"If further observations by Chandra and other satellites push the surface magnetic field limit lower, then theorists may have to dig deeper for an explanation of this enigmatic object," said Dr Nanda Rea, Institut de Ciències de l'Espai (ICE-CSIC, IEEC) in Barcelona, who led the discovery.

**More information:** 'A low-magnetic-field Soft Gamma Repeater' is published in the 14 October issue of *Science Express*.

Provided by University College London

Citation: Mysterious pulsar with hidden powers discovered (2010, October 14) retrieved 20 March 2024 from <https://phys.org/news/2010-10-mysterious-pulsar-hidden-powers.html>

|   |
|---|
| This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only. |
|---|