


# Planet warming will affect storms differently in Northern and Southern hemispheres

October 25 2010, by Morgan Bettex

---



Weather systems in the Southern and Northern hemispheres will respond differently to global warming, according to an MIT atmospheric scientist's analysis that suggests the warming of the planet will affect the availability of energy to fuel extratropical storms, or large-scale weather systems that occur at Earth's middle latitudes. The resulting changes will depend on the hemisphere and season, the study found.

More intense storms will occur in the [Southern Hemisphere](#) throughout the year, whereas in the Northern Hemisphere, the change in storminess will depend on the season — with more intense storms occurring in the winter and weaker storms in the summer. The responses are different because even though the [atmosphere](#) will get warmer and more humid due to [global warming](#), not all of the increased energy of the atmosphere will be available to power extratropical storms. It turns out that the

changes in available energy depend on the hemisphere and season, according to the study, published Monday in the *Proceedings of the National Academy of Sciences*.

Fewer extratropical storms during the summer in the Northern Hemisphere could lead to increased air pollution, as "there would be less movement of air to prevent the buildup of pollutants in the atmosphere," says author Paul O'Gorman, the Victor P. Starr Career Development Assistant Professor of Atmospheric Science in MIT's Department of Earth, Atmospheric and Planetary Sciences. Likewise, stronger storms year-round in the Southern Hemisphere would lead to stronger winds over the Antarctic Ocean, which would impact ocean circulation. Because the ocean circulation redistributes heat throughout the world's oceans, any change could impact the global climate.

O'Gorman's analysis examined the relationship between storm intensity and the amount of energy available to create the strong winds that fuel extratropical storms. After analyzing data compiled between 1981 and 2000 on winds in the atmosphere, he noticed that the energy available for storms depended on the season. Specifically, it increased during the winter, when extratropical storms are strong, and decreased during the summer, when they are weak.

Because this relationship could be observed in the current climate, O'Gorman was confident that available energy would be useful in relating temperature and storminess changes in global-warming simulations for the 21st century. After analyzing these simulations, he observed that changes in the energy available for storms were linked to changes in temperature and storm intensity, which depended on the season and hemisphere. He found that available energy increased throughout the year for the Southern Hemisphere, which led to more intense storms. But for the Northern Hemisphere, O'Gorman observed that available energy increased during the winter and decreased during

the summer.

This makes sense, O'Gorman says, because the changes in the strength of extratropical storms depend on where in the atmosphere the greatest warming occurs; if the warming is greatest in the lower part of the atmosphere, this tends to create stronger storms, but if it is greatest higher up, this leads to weaker storms. During the Northern Hemisphere summer, the warming is greatest at higher altitudes, which stabilizes the atmosphere and leads to less intense storms.

Although the analysis suggests that global warming will result in weaker Northern Hemisphere storms during the summer, O'Gorman says that it's difficult to determine the degree to which those storms will weaken. That depends on the interaction between the atmosphere and the oceans, and for the [Northern Hemisphere](#), this interaction is linked to how quickly the Arctic Ocean ice disappears. Unfortunately, climate scientists don't yet know the long-term rate of melting.

**More information:** "Understanding the varied response of the extratropical storm tracks to climate change," by Paul A. O'Gorman. Proceedings of the National Academy of Sciences, 25 October, 2010

Provided by Massachusetts Institute of Technology

Citation: Planet warming will affect storms differently in Northern and Southern hemispheres (2010, October 25) retrieved 3 May 2024 from <https://phys.org/news/2010-10-planet-affect-storms-differently-northern.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--