


Antihydrogen trapped for first time (w/ Video)

November 17 2010, By Robert Sanders



An octupole magnet was critical to trapping antihydrogen atoms by using their small magnetic moments. This simplified version shows how the north and south poles of strategically arranged magnets can immobilize a neutral antihydrogen atom that has a magnetic moment equivalent to a tiny bar magnet. Credit: Katie Bertsche

(PhysOrg.com) -- In the movie *Angels and Demons*, scientists have solved one of the most perplexing scientific problems: the capture and storage of antimatter. In real life, trapping atomic antimatter has never been accomplished, until now.

Physicists working at the European Organization for Nuclear Research

(CERN) in Geneva, Switzerland, have succeeded in trapping antihydrogen - the antimatter equivalent of the hydrogen atom - a milestone that could soon lead to experiments on a form of matter that disappeared mysteriously shortly after the birth of the universe 14 billion years ago.

The first artificially produced low energy antihydrogen atoms - consisting of a positron, or antimatter electron, orbiting an antiproton nucleus - were created at CERN in 2002, but until now the atoms have struck normal matter and annihilated in a flash of [gamma-rays](#) within microseconds of creation.

The ALPHA (Antihydrogen Laser PHysics Apparatus) experiment, an international collaboration that includes physicists from the University of California, Berkeley, and Lawrence Berkeley National Laboratory (LBNL), has now trapped 38 antihydrogen atoms, each for more than one-tenth of a second.

While the number and lifetime are insufficient to threaten the Vatican - in the 2000 novel and 2009 movie "Angels & Demons," a hidden vat of potentially explosive antihydrogen was buried under St. Peter's Basilica in Rome - it is a starting point for learning new physics, the researchers said.

"We are getting close to the point at which we can do some classes of experiments on the properties of antihydrogen," said Joel Fajans, UC Berkeley professor of physics, LBNL faculty scientist and ALPHA team member. "Initially, these will be crude experiments to test CPT symmetry, but since no one has been able to make these types of measurements on antimatter atoms at all, it's a good start."

CPT (charge-parity-time) symmetry is the hypothesis that physical interactions look the same if you flip the charge of all particles, change

their parity - that is, invert their coordinates in space - and reverse time. Any differences between antihydrogen and hydrogen, such as differences in their atomic spectrum, automatically violate CPT, overthrow today's "standard model" of particles and their interactions, and may explain why antimatter, created in equal amounts during the universe's birth, is largely absent today.


The team's results will be published online Nov. 17 in advance of its print appearance in the British journal *Nature*.

Antimatter, first predicted by physicist Paul Dirac in 1931, has the opposite charge of normal matter and annihilates completely in a flash of energy upon interaction with normal matter. While astronomers see no evidence of significant antimatter annihilation in space, [antimatter](#) is produced during high-energy particle interactions on earth and in some decays of radioactive elements. UC Berkeley physicists Emilio Segre and Owen Chamberlain created antiprotons in the Bevatron accelerator at the Lawrence Radiation Laboratory, now LBNL, in 1955, confirming their existence and earning the scientists the 1959 Nobel Prize in physics.

Slow antihydrogen was produced at CERN in 2002 thanks to an antiproton decelerator that slowed antiprotons enough for them to be used in experiments that combined them with a cloud of positrons. The ATHENA experiment, a broad international collaboration, reported the first detection of cold antihydrogen, with the rival ATRAP experiment close behind.

The ATHENA experiment closed down in 2004, to be superseded by ALPHA, coordinated by Jeffrey Hangst of the University of Aarhus in Denmark. Since then, the ALPHA and ATRAP teams have competed to trap antihydrogen for experiments, in particular, laser experiments to measure the antihydrogen spectrum (the color with which it glows) - and

gravity measurements. Before the recent results, the [CERN](#) experiments have produced - only fleetingly - tens of millions of antihydrogen atoms, Fajans said.



Top, a diagram of the region where antihydrogen is synthesized and trapped in ALPHA. Bottom a diagram of the electric potential in the trap region. Credit Nature

ALPHA's approach was to cool antiprotons and compress them into a matchstick-size cloud (20 millimeters long and 1.4 millimeters in diameter). Then, using autoresonance, a technique developed by UC Berkeley visiting professor Lazar Friedland and first explored in plasmas by Fajans and former U.C Berkeley graduate student Erik Gilson, the cloud of cold, compressed antiprotons is nudged to overlap a like-size [positron](#) cloud, where the two particles mate to form antihydrogen.

All this happens inside a magnetic bottle that traps the antihydrogen atoms. The magnetic trap is a specially configured magnetic field that Fajans and then-UC Berkeley undergraduate Andrea Schmidt first proposed, using an unusual and expensive octupole superconducting magnet to create a more stable plasma.


"For the moment, we keep antihydrogen atoms around for at least 172 milliseconds - about a sixth of a second - long enough to make sure we have trapped them," said colleague Jonathan Wurtele, UC Berkeley professor of physics and LBNL faculty scientist. Wurtele collaborated with LBNL visitor Katia Gomberoff, staff members Alex Friedman, David Grote and Jean-Luc Vay and with Fajans to simulate the new and original magnetic configurations.

Trapping antihydrogen isn't easy, Fajans said, because it is a neutral, or chargeless, particle. Magnetic bottles are generally used to trap charged particles, such as ionized atoms. These charged particles spiral along magnetic field lines until they encounter an electric field that bounces them back towards the center of the bottle.

Neutral antihydrogen, however, would normally be unaffected by these fields. But the team takes advantage of the tiny magnetic moment of the antihydrogen atom to trap it using a steeply increasing field - a so-called magnetic mirror - that reflects them backward toward the center. Because the magnetic moment is so small, the antihydrogen has to be very cold: less than about one-half degree above absolute zero (0.5 Kelvin). That means the team had to slow down the antiprotons by a factor of one hundred billion from their initial energy emerging from the antiproton decelerator.

Once trapped, the experimenters sweep out the lingering antiprotons with an electric field, then shut off the mirror fields and let the trapped antihydrogen atoms annihilate with normal matter. Surrounding detectors are sensitive to the charged pions that result from the proton-antiproton annihilation. Cosmic rays can also set off the detector, but their straight-line tracks can be easily distinguished, Fajans said. A few antiprotons could potentially remain in the trap, and their annihilations would look similar to those of antihydrogen, but the physicists' simulations show that such events can also be successfully distinguished

from antihydrogen annihilations.



The positions of the 38 real anti-atom annihilations (circles and triangles) match predicted antihydrogen distribution (gray dots in upper panel) but not the simulated distribution of bare antiprotons (colored dots in lower panel). Charged bare antiprotons would be steered to different clusters by different electric fields (red right bias, blue left bias, green no bias), but anti-atoms are neutral so their distribution is unaffected. (The violet star is an energetic positron.) Credit: courtesy of ALPHA team at CERN

During August and September of 2010, the team detected an antihydrogen atom in 38 of the 335 cycles of antiproton injection. Given that their detector efficiency is about 50 percent, the team calculated that it captured approximately 80 of the several million antihydrogen atoms produced during these cycles. Experiments in 2009 turned up six candidate antihydrogen atoms, but they have not been confirmed.

ALPHA continues to detect antihydrogen [atoms](#) at an increasing rate as the experimenters learn how to better tune their experiment, Fajans said.

Learning from antimatter

Before 1928, when anti-electrons were predicted on theoretical grounds by Paul Dirac, the existence of antimatter was unsuspected. In 1932 anti-electrons (positrons) were found in cosmic ray debris by Carl Anderson. The first antiprotons were deliberately created in 1955 at Berkeley Lab's Bevatron, the highest-energy particle accelerator of its day.

At first physicists saw no reason why antimatter and matter shouldn't behave symmetrically, that is, obey the laws of physics in the same way. But if so, equal amounts of each would have been made in the big bang in which case they should have mutually annihilated, leaving nothing behind. And if somehow that fate were avoided, equal amounts of matter and antimatter should remain today, which is clearly not the case.

In the 1960s, physicists discovered subatomic particles that decayed in a way only possible if the symmetry known as charge conjugation and parity (CP) had been violated in the process. As a result, the researchers realized, antimatter must behave slightly differently from ordinary matter. Still, even though some antiparticles violate CP, antiparticles moving backward in time ought to obey the same laws of physics as do ordinary particles moving forward in time. CPT symmetry (T is for time) should not be violated.

One way to test this assumption would be to compare the energy levels of ordinary electrons orbiting an ordinary proton to the energy levels of positrons orbiting an antiproton, that is, compare the spectra of ordinary hydrogen and antihydrogen atoms. Testing CPT symmetry with antihydrogen atoms is a major goal of the ALPHA experiment.

More information: *Nature* paper: [www.nature.com/nature/journal/...ull/nature09610.html](http://www.nature.com/nature/journal/full/nature09610.html)

Provided by University of California - Berkeley

Citation: Antihydrogen trapped for first time (w/ Video) (2010, November 17) retrieved 6 February 2023 from <https://phys.org/news/2010-11-antihydrogen.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.