

# Extreme global warming in the ancient past

November 10 2010

---



The image shows the the scientific drilling ship JOIDES Resolution docked in Hobart, Tasmania. Credit: John Beck, IODP

Variations in atmosphere carbon dioxide around 40 million years ago were tightly coupled to changes in global temperature, according to new findings published in the journal *Science*. The study was led by scientists at Utrecht University, working with colleagues at the NIOZ Royal Netherlands Institute for Sea Research and the University of Southampton.

"Understanding the relationship between the Earth's climate and [atmospheric carbon dioxide](#) in the geological past can provide insight into the extent of future global warming expected to result from [carbon dioxide](#) emission caused by the activities of humans," said Dr Steven Bohaty of the University of Southampton's School of Ocean and Earth Science (SOES) based at the National Oceanography Centre in

Southampton.

It has been known for some time that the long-term warmth of the Eocene (~56 to 34 million years ago) was associated with relatively high atmospheric carbon dioxide levels. However, scientists were previously unable to demonstrate tight-coupling between variations in atmospheric carbon dioxide and shorter-term changes in global climate.

To fill this gap in knowledge, the authors of the new study focused on one of the hottest episodes of Earth's [climate history](#) – the Middle Eocene Climatic Optimum (MECO), which occurred around 40 million years ago.

Algae use photosynthesis to harvest the energy of the sun, converting carbon dioxide and water into the organic molecules required for growth. Different isotopes of carbon are incorporated into these molecules depending on the environmental conditions under which algae grow. Ancient climate can therefore be reconstructed by analysing the carbon isotope ratios of molecules preserved in fossilised algae.

The researchers took this approach to reconstruct variations in carbon dioxide levels across the MECO warming event, using fossilised algae preserved in sediment cores extracted from the seafloor near Tasmania, Australia, by the Ocean Drilling Program. They refined their estimates of carbon dioxide levels using information on the past marine ecosystem derived from studying changes in the abundance of different groups of fossil plankton.

Their analyses indicate that MECO carbon dioxide levels must have at least doubled over a period of around 400,000 years. In conjunction with these findings, analyses using two independent molecular proxies for sea surface temperature show that the climate warmed by between 4 and 6 degrees Celsius over the same period.

"We found a close correspondence between carbon dioxide levels and sea surface temperature over the whole period, suggesting that increased amounts of carbon dioxide in the atmosphere played a major role in global warming during the MECO," said Bohaty.

The researchers consider it likely that elevated atmospheric carbon dioxide levels during the MECO resulted in increased global temperatures, rather than vice versa, arguing that the increase in carbon dioxide played the lead role.

"The change in carbon dioxide 40 million years ago was too large to have been the result of temperature change and associated feedbacks," said co-lead author Peter Bijl of Utrecht University. "Such a large change in carbon dioxide certainly provides a plausible explanation for the changes in Earth's temperature."

The researchers point out that the large increase in atmospheric carbon dioxide indicated by their analysis would have required a natural carbon source capable of injecting vast amounts of carbon into the atmosphere.

The rapid increase in atmospheric carbon dioxide levels around 40 million years ago approximately coincides with the rise of the Himalayas and may be related to the disappearance of an ocean between India and Asia as a result of plate tectonics – the large scale movements of the Earth's rocky shell (lithosphere). But, as explained by Professor Paul Pearson of Cardiff University in a perspective article accompanying the *Science* paper, the hunt is now on to discover the exact cause.

**More information:**

-- Publication: Bijl, P. K., Houben, A. J. P., Schouten, S., Bohaty, S. M., Sluijs, A., Reichert, G-J., Sinninghe Damsté, J. S. & Brinkhuis, H. Transient middle Eocene atmospheric CO<sub>2</sub> and temperature variations. *Science* 330, 819 - 8215 (2010). [DOI: 10.1126/science.1193654](https://doi.org/10.1126/science.1193654)

-- Science Perspective: Pearson, P. N. Increased atmospheric CO<sub>2</sub> during the middle Eocene. *Science* 330, 763-764 (2010). [DOI: 10.1126/science.1197894](https://doi.org/10.1126/science.1197894)

Provided by National Oceanography Centre, Southampton

Citation: Extreme global warming in the ancient past (2010, November 10) retrieved 25 April 2024 from <https://phys.org/news/2010-11-extreme-global-ancient.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.