

The physics of coffee rings

November 24 2010

For centuries, intellectuals have met at the ring-stained surfaces of coffee shops to pore over the most pressing problems of the day -- but has anyone ever pondered the coffee rings they left behind? What causes the formation of stain patterns left behind by coffee droplets on a surface?

You might think [coffee](#) ring formation, first described quantitatively by Deegan et al in a heavily cited article, is the most widely and ritualistically performed experiment in the world, given the prevalence of caffeine in cultures. But most of us lack the [scanning electron microscope](#) and mathematical models to evaluate our stain data properly, or reach meaningful conclusions beyond "Use a coaster."

Now Shreyas Mandre of Brown University, Ning Wu from Colorado School of Mines and L. Mahadevan and Joanna Aizenberg from Harvard University have devised a [predictive model](#) that combines laboratory studies of microscopic glass particles in solution with mathematical theories to predict the existence, thickness and length of the banded ring patterns that formed.

Their results, presented today at the American Physical Society Division of Fluid Dynamics meeting in Long Beach, CA, suggest the patterned deposition of particles can be controlled by altering physical parameters such as evaporation and [surface tension](#) -- and perhaps one day manipulated to create small-particle tools.

"Controlling the ring deposition process would be useful for creating

such things as new microphysics tools operating at a scale where pliers or other traditional tools for moving particles cannot operate," notes Mandre.

The team found that during ring deposition, a particle layer of uniform thickness is deposited if the concentration is above a certain threshold. Below that threshold the deposits form non-uniform bands. The threshold is formed because evaporation at the solid-liquid interface of the rim occurs faster than a replenishing flow of water from the center of the droplet can replace the evaporating rim fluid. This leaves the particles on the rim high, dry -- and deposited.

Exploiting this competition between evaporation and replenishment is the key to controlling the process as a microtool, says Mandre. Potential applications include printing, making industrial coatings, fabricating electronics, and designing new medicines.

More information: The presentation, "Coffee ring deposition in bands" is on Tuesday, November 23, 2010. Abstract: meetings.aps.org/Meeting/DFD10/Event/134397

Provided by American Institute of Physics

Citation: The physics of coffee rings (2010, November 24) retrieved 18 April 2024 from <https://phys.org/news/2010-11-physics-coffee.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--