

Predatory bugs can save cornfields

November 30 2010

One of the worst pests of corn in the world, the corn rootworm, may owe its worldwide success partly to its larvae's nasty, sticky blood.

U.S. Department of Agriculture (USDA) [entomologist](#) Jonathan G. Lundgren and his colleagues discovered this recently, working with CABI researchers in Delémont, Switzerland, and Hódmezővásárhely, Hungary. The discovery could lead to development of ways to overcome these defenses as part of sustainable, ecologically based pest management methods.

Lundgren works at the Agricultural Research Service (ARS) North Central Agricultural Research Laboratory in Brookings, S.D. ARS is USDA's principal intramural scientific research agency, and this research supports the USDA priority of promoting international food security. CABI is an international not-for-profit organization that researches natural ways of controlling [pests](#), and they have been helping to lead the effort against corn rootworm's European invasion.

The experiments with CABI are the latest in Lundgren's research on [corn](#) rootworm predators. Although rootworms have been a major pest for 100 years, this is remarkably the first comprehensive research program on [corn rootworm](#) predators to be conducted.

In lab and field experiments in the United States and abroad, the rootworm larvae's sticky blood caused certain species of predators to quickly back off. The foul-tasting blood coagulated in the predators' mouths, temporarily gluing them shut. Predators repelled by the

rootworm larvae's blood included ground beetles and ants.

Wolf spiders, on the other hand, had a hearty appetite for rootworms. When insects such as spiders suck fluids from prey rather than chewing their victims, they may be able to bypass the ability of the blood to stick and linger.

The experiments with CABI involved two years of lab and field experiments, begun in 2007, in the United States and Hungary. In the Brookings laboratory, Lundgren and colleagues offered hungry predators a smorgasbord of rootworm [larvae](#) and pupae. In all, they have tested 10 different predator species from Europe and North America.

The results have led Lundgren to research managing crop fields to encourage large and diverse predator populations.

More information: Papers on this research have been published recently in Biocontrol Science and Technology, Ecological Applications, and the Journal of Applied Entomology. Also, a paper is scheduled for publication in the Environmental Entomology journal.

Provided by United States Department of Agriculture

Citation: Predatory bugs can save cornfields (2010, November 30) retrieved 25 April 2024 from <https://phys.org/news/2010-11-predatory-bugs-cornfields.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.