

# New property in warm superconductors discovered

November 17 2010

---

(PhysOrg.com) -- Led by Simon Fraser University physicist Jeff Sonier, scientists at TRIUMF have discovered something that they think may severely hinder the creation of room-temperature (37 degrees Celsius) superconductors.

For 25 years, they've speculated that magnetism could be a problem.

The *Proceedings of the National Academy of Sciences* has published the finding that there is a weak magnetism in a certain type of lanthanum-based copper oxide material, which is the closest known warm-temperature superconductor.

Sonier, the lead scientist in this research, says: "The search for room-temperature [superconductivity](#) is big news. The cover story of the June 2010 issue of *Scientific American* predicted the discovery would be one of the '12 events that will change everything.'"

Superconductors, materials that have zero electrical resistance, could potentially drive ever day devices in electronics, medicine and transportation, but are super expensive because they only operate at extremely low temperatures. If [superconductors](#) were operational at room temperature they wouldn't need to be driven by expensive cooling systems using liquid helium.

When charge carriers are added to copper oxide materials, known as cuprates, they are capable of superconductivity. Some cuprates function

at -140 degrees Celsius, a temperature markedly above -240 degrees Celsius, which is the normal operational temperature of all other kinds of superconducting materials.

Adding charge carriers (electric charge carrying particle) is known as chemical doping. With increased chemical doping the operational temperature of a cuprate superconductor rises to a certain point and then collapses.

Until this latest research, scientists could only speculate on whether a competing magnetic phase might exist during high chemical doping and ultimately destroy their superconductivity.

Sonier and his colleagues used a subatomic particle, called a muon, to microscopically probe the magnetic nature of a cuprate. This led them to discover that a strange kind of magnetism appears to accompany the destruction of superconductivity during high chemical doping.

The scientists are now trying to figure out the origin of the magnetism and whether it actually competes with superconductivity.

Sonier says, “Understanding what destroys superconductivity during high chemical doping could provide a vital clue about the microscopic mechanism responsible for high-temperature superconductivity. Knowledge of this would be a monumental step toward making a room-temperature superconductor.”

**More information:** Direct search for a ferromagnetic phase in a heavily overdoped nonsuperconducting copper oxide, *PNAS* October 5, 2010 vol. 107 no. 40 17131-17134, [doi:10.1073/pnas.1007079107](https://doi.org/10.1073/pnas.1007079107)

Provided by Simon Fraser University

Citation: New property in warm superconductors discovered (2010, November 17) retrieved 20 April 2024 from <https://phys.org/news/2010-11-property-superconductors.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.