

Pulsating star mystery solved

November 24 2010



This artist's impression shows the double star OGLE-LMC-CEP0227 in our neighboring galaxy the Large Magellanic Cloud. The smaller of the two stars is a pulsating Cepheid variable and the orientation of the system is such that the stars eclipse each other during their orbits. Studies of this very rare system have allowed astronomers to measure the Cepheid mass with unprecedented accuracy. Credit: ESO

(PhysOrg.com) -- By discovering the first double star where a pulsating Cepheid variable and another star pass in front of one another, an international team of astronomers has solved a decades-old mystery. The rare alignment of the orbits of the two stars in the double star system has allowed a measurement of the Cepheid mass with unprecedented accuracy. The new result shows that the prediction from stellar pulsation theory is spot on, while the prediction from stellar evolution theory is at odds with the new observations.

The new results, from a team led by Grzegorz Pietrzynski (Universidad

de Concepción, Chile, Obserwatorium Astronomiczne Uniwersytetu Warszawskiego, Poland), appear in the 25 November 2010 edition of the journal *Nature*.

Grzegorz Pietrzynski introduces this remarkable result: "By using the HARPS instrument on the 3.6-metre telescope at ESO's La Silla Observatory in Chile, along with other telescopes, we have measured the mass of a Cepheid with an accuracy far greater than any earlier estimates. This new result allows us to immediately see which of the two competing theories predicting the masses of Cepheids is correct."

Classical Cepheid Variables, usually called just Cepheids, are unstable stars that are larger and much brighter than the Sun. They expand and contract in a regular way, taking anything from a few days to months to complete the cycle. The time taken to brighten and grow fainter again is longer for stars that are more luminous and shorter for the dimmer ones. This remarkably precise relationship makes the study of Cepheids one of the most effective ways to measure the distances to nearby galaxies and from there to map out the scale of the whole Universe.

Unfortunately, despite their importance, Cepheids are not fully understood. Predictions of their masses derived from the theory of pulsating stars are 20% less than predictions from the theory of the evolution of stars. This embarrassing discrepancy has been known since the 1960s.

To resolve this mystery, astronomers needed to find a double star containing a Cepheid where the orbit happened to be seen edge-on from Earth. In these cases, known as eclipsing binaries, the brightness of the two stars dims as one component passes in front of the other, and again when it passes behind the other star. In such pairs [astronomers](#) can determine the masses of the stars to high accuracy. Unfortunately neither Cepheids nor eclipsing binaries are common, so the chance of finding

such an unusual pair seemed very low. None are known in the Milky Way.



This wide field view of part of our neighbouring galaxy, the Large Magellanic Cloud, was created from photographs taken in red and blue light as part of the Digitized Sky Survey 2. The remarkable double star OGLE-LMC-CEP0227 lies at the centre of the picture, just one of huge numbers of faint stars. Studies of this very rare system have allowed astronomers to measure mass of a Cepheid variable star with unprecedented accuracy. The field of view is approximately 2.8 degrees across. Credit: ESO

Wolfgang Gieren, another member of the team, takes up the story: "Very recently we actually found the double star system we had hoped for among the stars of the Large Magellanic Cloud. It contains a Cepheid variable star pulsating every 3.8 days. The other star is slightly bigger and cooler, and the two stars orbit each other in 310 days. The true binary nature of the object was immediately confirmed when we observed it with the HARPS spectrograph on La Silla."

The observers carefully measured the brightness variations of this rare object, known as OGLE-LMC-CEP0227, as the two stars orbited and passed in front of one another. They also used HARPS and other spectrographs to measure the motions of the stars towards and away from the Earth — both the orbital motion of both stars and the in-and-out motion of the surface of the Cepheid as it swelled and contracted.

This very complete and detailed data allowed the observers to determine the orbital motion, sizes and masses of the two stars with very high accuracy — far surpassing what had been done before for a Cepheid. The mass of the Cepheid is now known to about 1% and agrees exactly with predictions from the theory of stellar pulsation. However, the larger mass predicted by stellar evolution theory was shown to be significantly in error.

The much-improved mass estimate is only one outcome of this work, and the team hopes to find other examples of these remarkably useful pairs of [stars](#) to exploit the method further. They also believe that from such binary systems they will eventually be able to pin down the distance to the Large Magellanic Cloud to 1%, which would mean an extremely important improvement of the cosmic distance scale.

More information: This research was presented in a paper to appear in the journal *Nature* on 25 November 2010.

Provided by ESO

Citation: Pulsating star mystery solved (2010, November 24) retrieved 28 April 2024 from <https://phys.org/news/2010-11-pulsating-star-mystery.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.