


# Saturn then and now: 30 years since Voyager visit

November 12 2010, By Jia-Rui C. Cook

---




The day NASA's Voyager 1 spacecraft made its closest approach to Saturn, it revealed the kinks in one of Saturn's narrowest rings (left). Images from the Cassini spacecraft (right) finally allowed scientists to understand how Saturn's moons Prometheus and Pandora sculpt the ring's kinked shape.

Ed Stone, project scientist for NASA's Voyager mission, remembers the first time he saw the kinks in one of Saturn's narrowest rings. It was the day the Voyager 1 spacecraft made its closest approach to the giant ringed planet, 30 years ago. Scientists were gathering in front of television monitors and in one another's offices every day during this heady period to pore over the bewildering images and other data

streaming down to NASA's Jet Propulsion Laboratory in Pasadena, Calif.

Stone drew a crude sketch of this scalloped, multi-stranded ring, known as the F ring, in his notebook, but with no explanation next to it. The innumerable particles comprising the broad rings are in near-circular orbits about Saturn. So, it was a surprise to find that the F ring, discovered just a year before by NASA's Pioneer 11 [spacecraft](#), had clumps and wayward kinks. What could have created such a pattern?

"It was clear Voyager was showing us something different at Saturn," said Stone, now based at the California Institute of Technology in Pasadena. "Over and over, the spacecraft revealed so many unexpected things that it often took days, months and even years to figure them out."




The puzzling hexagon-shaped weather pattern around Saturn's north pole was first found in images from Voyager 2 (left). Cassini has obtained higher-resolution pictures of the hexagon (such as the one on the right, taken in 2009) - which tell scientists the hexagon is a remarkably stable wave in one of the jet streams that remains 30 years later.

The F ring curiosity was only one of many strange phenomena discovered in the Voyager close encounters with Saturn, which occurred on Nov. 12, 1980, for Voyager 1, and Aug. 25, 1981, for [Voyager 2](#). The Voyager encounters were responsible for finding six small moons and revealing the half-young, half-old terrain of Enceladus that had to point to some kind of geological activity.

Images from the two encounters also exposed individual storms roiling the planet's atmosphere, which did not show up at all in data from Earth-based telescopes. Scientists used Voyager data to resolve a debate about whether Titan had a thick or thin atmosphere, finding that Titan was shrouded in a thick haze of hydrocarbons in a nitrogen-rich atmosphere. The finding led scientists to predict there could be seas of liquid methane and ethane on Titan's surface.

"When I look back, I realize how little we actually knew about the solar system before Voyager," Stone added. "We discovered things we didn't know were there to be discovered, time after time."



Scientists first saw these transient clouds of tiny particles known as "spokes" in images from NASA's Voyager spacecraft. An electrostatic charge appears to be levitating tiny ring particles above the ring plane, but scientists are still figuring out how the particles get that charge as they analyze images from NASA's Cassini spacecraft.

In fact, the Voyager encounters sparked so many new questions that another spacecraft, NASA's Cassini, was sent to probe those mysteries. While Voyager 1 got to within about 126,000 kilometers (78,300 miles) above Saturn's cloud tops, and Voyager 2 approached as close as about 100,800 kilometers (62,600 miles), Cassini has dipped to this altitude and somewhat lower in its orbits around Saturn since 2004.

Because of Cassini's extended journey around Saturn, scientists have found explanations for many of the mysteries first seen by Voyager. Cassini has uncovered a mechanism to explain the new terrain on [Enceladus](#) – tiger stripe fissures with jets of water vapor and organic particles. It revealed that Titan indeed does have stable lakes of liquid hydrocarbons on its surface and showed just how similar to Earth that

moon really is. Data from Cassini have also resolved how two small moons discovered by Voyager – Prometheus and Pandora – tug on the F ring to create its kinked shape and wakes that form snowballs.

"Cassini is indebted to Voyager for its many fascinating discoveries and for paving the way for Cassini," said Linda Spilker, Cassini project scientist at JPL, who started her career working on Voyager from 1977 to 1989. "On Cassini, we still compare our data to Voyager's and proudly build on Voyager's heritage."

But Voyager left a few mysteries that Cassini has not yet solved. For instance, scientists first spotted a hexagonal weather pattern when they stitched together Voyager images of Saturn's north pole. Cassini has obtained higher-resolution pictures of the hexagon – which tells scientists it's a remarkably stable wave in one of the jet streams that remains 30 years later – but scientists are still not sure what forces maintain the hexagon.

Even more perplexing are the somewhat wedge-shaped, transient clouds of tiny particles that Voyager discovered orbiting in Saturn's B ring. Scientists dubbed them "spokes" because they looked like bicycle spokes. Cassini scientists have been searching for them since the spacecraft first arrived. As [Saturn](#) approached equinox, and the sun's light hit the rings edge-on, the spokes did reappear in the outer part of Saturn's B ring. But Cassini scientists are still testing their theories of what might be causing these odd features.

"The fact that we still have mysteries today goes to show how much we still have to learn about our solar system," said Suzanne Dodd, Voyager's project manager, based at JPL. "Today, the Voyager spacecraft continue as pioneers traveling toward the edge of our solar system. We can't wait for the Voyager spacecraft to enter interstellar space – true outer space – and make more unexpected discoveries."

[Voyager 1](#), which was launched on Sept. 5, 1977, is currently about 17 billion kilometers (11 billion miles) away from the sun. It is the most distant spacecraft. Voyager 2, which was launched on Aug. 20, 1977, is currently about 14 billion kilometers (9 billion miles) away from the sun.

The Voyagers were built by JPL, which continues to operate both spacecraft. Caltech manages JPL for NASA. The Cassini-Huygens mission is a cooperative project of NASA, the European Space Agency and the Italian Space Agency. JPL manages Cassini for NASA. The Cassini orbiter and its two onboard cameras were designed, developed and assembled at JPL.

Provided by JPL/NASA

Citation: Saturn then and now: 30 years since Voyager visit (2010, November 12) retrieved 2 May 2024 from <https://phys.org/news/2010-11-saturn-years-voyager.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--