


Vacuum arcs spark new interest

November 8 2010



Even in a vacuum, voltages can literally tear metal apart. Current research is using molecular dynamics to show what happens at the atomic level when material is damaged by arcing in vacuum. Credit: J. Norem, Z. Insepov, Th. Proslir, D. Huang, S. Mahalingam, and S. Veitzer

Whenever two pieces of metal at different voltages are brought near each other, as when an appliance is plugged into a live socket, there is a chance there will be an arc between them. Most of the arcs people see are a breakdown of the gas between the metal surfaces, but this type of breakdown can also occur in a vacuum. This vacuum breakdown, which until recently has not been well understood, has implications for applications from particle accelerators to fusion reactors.

As part of an effort to understand the maximum accelerating field in particle accelerators, scientists at Argonne National Laboratory have been modeling the processes involved in vacuum breakdown. Now, a new model of this phenomenon is beginning to reveal what is happening

in these arcs, and scientists are studying a number of new phenomena associated with them.

In this new model, the breakdown arc is triggered by the electric field in the [vacuum](#) gap literally tearing the metal apart. (The same force that causes "static cling" can be very powerful for high electric fields, particularly at tiny corners, and in cracks where the fields are intensified by the local geometry of the surface.) After the metal is torn apart, the fragments should become ionized and form microscopic plasmas that are very dense and cold (for a plasma). Because of the high densities in these plasmas, the surface fields inside the arc quickly become even stronger than they were at breakdown. The arc becomes very damaging to the metal surface over a comparatively large area, eventually leaving a pit that should be visible to the naked eye.

While this model seems to be internally consistent, researchers want to use it to produce predictions that can be verified experimentally. Current research is using [molecular dynamics](#) to show what happens at the [atomic level](#) when the material is torn apart, plasma modeling codes to show how the plasma initially forms and what its properties are, and electrohydrodynamics to show how the surface of the arc pit is affected. While current results seem generally consistent with existing experimental measurements, more precise tests are being developed.

In principle, a better understanding of the precise causes of electrical breakdown should suggest solutions that are relevant to fusion reactors and space communications, among other things.

Provided by American Physical Society

Citation: Vacuum arcs spark new interest (2010, November 8) retrieved 26 April 2024 from <https://phys.org/news/2010-11-vacuum-arcs.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.