


Dark matter could transfer energy in the Sun

December 1 2010



This is the Sun, picked up from the space telescope Hinode X-ray. Credit: Hinode JAXA/NASA/PPARC

Researchers from the Institute for Corpuscular Physics (IFIC) and other European groups have studied the effects of the presence of dark matter in the Sun. According to their calculations, low mass dark matter particles could be transferring energy from the core to the external parts of the Sun, which would affect the quantity of neutrinos that reach the Earth.

"We assume that the [dark matter particles](#) interact weakly with the Sun's atoms, and what we have done is calculate at what level these

interactions can occur, in order to better describe the structure and evolution of the Sun", Marco Taoso, researcher at the IFIC, a combined centre of the Spanish National Research Council and the University of Valencia, explains to SINC.

The astrophysical observations suggest that our galaxy is situated in a halo of dark matter particles. According to the models, some of these particles, the WIMPs (Weakly Interacting Massive Particles) interact weakly with other normal ones, such as atoms, and could be building up on the inside of stars. The study, recently published in the journal *Physical Review D*, carries out an in-depth study of the case of the Sun in particular.

"When the WIMPs pass through the Sun they can break up the atoms of our star and lose energy. This prevents them from escaping the gravitational force of the Sun which captures them, and they become trapped, orbiting inside it, with no way of escaping", the researcher points out.

The dark matter cools down the Sun's core

Scientists believe that the majority of the dark matter particles gather together in the centre of the Sun, but in their elliptic orbits they also travel to the outer part, interacting and exchanging with the solar atoms. In this way, the WIMPs transport the energy from the burning central core to the cooler peripheral parts.

"This effect produces a cooling down of the core, the region from where the neutrinos originate due to the nuclear reactions of the Sun", Taoso points out. "And this corresponds to a reduction in the flux of solar neutrinos, since these depend greatly on the temperature of the core".

The neutrinos that reach the Earth can be measured by means of

different techniques. These data can be used to detect the modifications of the solar temperature caused by the WIMPs. The transport of energy by these particles depends on the likelihood of them interacting with the atoms, and the "size" of these interactions is related to the reduction in the neutrino flux.

"As a result, current data about solar neutrinos can be used to put limits on the extent of the interactions between dark matter and [atoms](#), and using numerical codes we have proved that certain values correspond to a reduction in the flux of solar neutrinos and clash with the measurements", the scientist reveals.

The team has applied their calculations to better understand the effects of low mass [dark matter](#) particles (between 4 and 10 giga-electronvolts). At this level we find models that attempt to explain the results of experiments such as DAMA (beneath an Italian mountain) or CoGent (in a mine in the USA), which look for dark material using "scintillators" or WIMP detectors.

Debate about WIMP and solar composition

This year another study by scientists from Oxford University (United Kingdom) also appeared. It states that WIMPs not only reduce the fluxes of solar [neutrinos](#), but also, furthermore, modify the structure of the Sun and can explain its composition.

"Our calculations, however, show that the modifications of the star's structure are too small to support this claim and that the [WIMPs](#) cannot explain the problem of the composition of the [sun](#)", Taoso concludes.

More information: Marco Taoso, Fabio Iocco, Georges Meynet, Gianfranco Bertone y Patrick Eggenberger. "Effect of low mass dark matter particles on the Sun". *Physical Review D* 82 (8), Oct 2010. [DOI](#):

[10.1103/PhysRevD.82.083509](https://arxiv.org/abs/1011.0835)

Provided by FECYT - Spanish Foundation for Science and Technology

Citation: Dark matter could transfer energy in the Sun (2010, December 1) retrieved 1 April 2023 from <https://phys.org/news/2010-12-dark-energy-sun.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.