

A study analyzes the movement of tree sap

December 13 2010



This picture show a regular plant which is used in the research about the movement of sap in order to apply these advances to new hydraulic systems or to suction pumps. Credit: UC3M

Spanish scientists at Carlos III University of Madrid use 3-D modeling to analyze the mechanisms used to by trees to transport water in their interior. The objective: to discover the keys to the movement of sap in order to apply these advances to new hydraulic systems or to suction pumps.

The researchers decided to embark on this study in order to find out which mechanisms are used by plants when they extract water from very dry or somewhat inhospitable land. "In the case of mangrove swamps, for example, the plants are able to extract freshwater from a saltwater



environment, despite the fact that the osmotic pressure should make quite the opposite happen", explains Professor José Luis Pérez Díaz, who studies this type of relatively unknown phenomenon as part of a new line of research that the Department of Mechanical Engineering at UC3M has begun.

The objective of the study is to learn what type of mechanism the plants use when extracting the water and transporting it from the roots to the leaves. To do this, the researchers have generated a model that represents the microscopic structure of the trunk of a European beech tree (Fagus sylvatica) in order to study the changes produced when the water moves through its interior; they have published some of the results of their research in a recent issue of the *Journal of Biological Physics*.



This is an infographic about the study to find out which mechanisms are used by plants when they extract water. Credit: Carlos III University of Madrid

The three-dimensional model created by these UC3M scientists allows them to analyze the transpiration and absorption of the bark of the tree, as well as the pressure in the different types of conduits, such as the floem or xylem, that transport fluids inside the trunk. The first is a



vascular tissue that carries sugars and other synthesized nutrients from the organs that produce them to the organs where they are consumed and stored, in both upward and downward directions. Xylem, on the other hand, is different type of tissue that is adapted especially for conveying water upward for the length of the plant; its elements are arranged in longitudinal rows, through more or less continuous sections of conduits.

The main conclusion of this study is that the <u>sap</u> in the trunks of <u>trees</u> is in a pressurized situation. It demonstrates, then, that when the pressure is positive in the conduits of the xylem as well as in those of the floem, the model expands in the radial direction. However, when the pressure is negative in the xylem and positive in the floem, which is what is believed to occur during the day, the model contracts in the radial direction. "Our results are not absolutely conclusive yet, but they set us on a path to continue our study and to find out more about these processes", says Professor Pérez Díaz.

This line of research could have interesting applications for hydraulic systems or devices for water extraction, for example. "Currently – the expert points out – there is no water suction pump capable of raising water more than ten meters at normal atmospheric pressure, but a sequoia tree can raise <u>water</u> to a height of 100 meters, which I think means that anything we can learn from <u>plants</u> is going to be of great interest to people working in this field", he concludes.

Provided by Carlos III University of Madrid

Citation: A study analyzes the movement of tree sap (2010, December 13) retrieved 21 May 2024 from <u>https://phys.org/news/2010-12-movement-tree-sap.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is



provided for information purposes only.