

Smithsonian instrument 'fills the gap,' views sun's innermost corona

January 4 2011



This photograph of the sun, taken by the Atmospheric Imaging Assembly instrument on NASA's Solar Dynamics Observatory, shows how image processing techniques developed at SAO can reveal the faint, inner corona. At the sun's limb, prominences larger than the Earth arc into space. Bright active regions like the one on the Sun's face at lower center are often the source of huge eruptions known as coronal mass ejections. Credit: NASA/LMSAL/SAO

During a total eclipse of the Sun, skywatchers are awed by the shimmering corona -- a faint glow that surrounds the Sun like gossamer flower petals. This outer layer of the Sun's atmosphere is, paradoxically, hotter than the Sun's surface, but so tenuous that its light is overwhelmed by the much brighter solar disk. The corona becomes visible only when



the Sun is blocked, which happens for just a few minutes during an eclipse.

Now, an instrument on board NASA's Solar Dynamics Observatory (SDO), developed by Smithsonian scientists, is giving unprecedented views of the innermost corona 24 hours a day, 7 days a week.

"We can follow the corona all the way down to the Sun's surface," said Leon Golub of the Harvard-Smithsonian Center for Astrophysics (CfA).

Previously, solar astronomers could observe the corona by physically blocking the solar disk with a coronagraph, much like holding your hand in front of your face while driving into the setting Sun. However, a <u>coronagraph</u> also blocks the area immediately surrounding the Sun, leaving only the outer corona visible.

The Atmospheric Imaging Assembly (AIA) instrument on SDO can "fill" this gap, allowing astronomers to study the corona all the way down to the Sun's surface. The resulting images highlight the ever-changing connections between gas captured by the Sun's magnetic field and gas escaping into interplanetary space.

The Sun's magnetic field molds and shapes the corona. Hot solar plasma streams outward in vast loops larger than Earth before plunging back onto the Sun's surface. Some of the loops expand and stretch bigger and bigger until they break, belching plasma outward.





This zoomed-in image shows how the sun's magnetic field shapes hot coronal plasma. Photos like this highlight the ever-changing connections between gas captured by the sun's magnetic field and gas escaping into interplanetary space. Credit: NASA/LMSAL/SAO

"The AIA solar images, with better-than-HD quality views, show <u>magnetic structures</u> and dynamics that we've never seen before on the Sun," said CfA astronomer Steven Cranmer. "This is a whole new area of study that's just beginning."

Cranmer and CfA colleague Alec Engell developed a computer program for processing the AIA images above the Sun's edge. These processed images imitate the blocking-out of the Sun that occurs during a total solar eclipse, revealing the highly dynamic nature of the inner <u>corona</u>. They will be used to study the initial eruption phase of coronal mass ejections (CMEs) as they leave the <u>Sun</u> and to test theories of solar wind acceleration based on magnetic reconnection.

Provided by Harvard-Smithsonian Center for Astrophysics

Citation: Smithsonian instrument 'fills the gap,' views sun's innermost corona (2011, January 4) retrieved 3 May 2024 from https://phys.org/news/2011-01-smithsonian-instrument-gap-views-sun.html



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.