

Wheat resistance genes failing, new approach needed to stop flies

January 24 2011

(PhysOrg.com) -- Many of the genes that allow wheat to ward off Hessian flies are no longer effective in the southeastern United States, and care should be taken to ensure that resistance genes that so far haven't been utilized in commercial wheat lines are used prudently, according to U.S. Department of Agriculture and Purdue University scientists.

An analysis of wheat lines carrying [resistance genes](#) from dozens of locations throughout the Southeast showed that some give little or no resistance to the Hessian fly, a major pest of wheat that can cause millions of dollars in damage to wheat crops each year. Others, even those considered the most effective, are allowing wheat to become susceptible to the fly larvae, which feed on and kill the plants.

Wheat resistance genes recognize avirulent Hessian flies and activate a defense response that kills the [fly larvae](#) attacking the plant. However, this leads to strains of the fly that can overcome [resistant wheat](#), much like insects becoming resistant to pesticides.

"The number of genes available to protect wheat is limited. There really aren't that many," said Richard Shukle, a research scientist with the USDA Agricultural Research Service [Crop Production](#) and Pest Control Research Unit and Purdue adjunct associate professor of entomology. "In the Southeast, having multiple generations of Hessian fly each year enhances the ability of these flies to overcome wheat's resistance."

Sue Cambron, a USDA Agricultural Research Service research support scientist, received Hessian flies from 20 locations in Alabama, Georgia, North Carolina, South Carolina and Louisiana and used them to infest 21 varieties of wheat that each contained different resistance genes, most of which have been deployed in commercial wheat, and a few that haven't yet. While the study did not include all of the 33 named resistance genes, it did show that only five of the 21 genes evaluated would provide effective resistance to flies in the Southeast, and none was effective in all the Southeast locations.

"Even some of the newer genes that haven't been deployed in cultivars weren't too effective," Cambron said.

That's because flies have likely interacted with, and adapted to, those genes already, said Brandi Schemerhorn, a USDA-ARS entomologist and Purdue assistant professor of entomology. She said it's possible that some of the genes were introduced to flies unintentionally in plots where wheat cultivars with those genes were being tested for suitability to Southeast climates. The resistance genes also could have come from other plants, such as rye, and the flies may already have started to overcome those genes.

Schemerhorn said she suspects a certain number of flies in any population have the ability to overcome any wheat resistance gene, which defends against the flies' ability to feed on the plant and starves the insect larvae. When a resistance gene kills off some of the flies, the survivors breed and eventually establish a population that renders the gene ineffective.

"We're creating a system in which the fly is becoming more virulent," Schemerhorn said. "What we have to do is slow down that adaptation or virulence."

Shukle and Schemerhorn suggest stacking genes in a wheat cultivar. There are only a few genes that haven't been deployed, and they believe combining two of those would be the best option.

"With a small number of identified resistance genes, we can't afford to release wheat lines with only one resistance gene," Shukle said. "If you deploy two different resistance genes, it's unlikely that a population of flies could overcome both of them."

Schemerhorn is working to combine two of the unreleased genes for testing with [Hessian fly](#) populations.

The USDA funded the research. Shukle, Schemerhorn and Cambron collaborated with David Buntin, a University of Georgia professor of entomology; Randy Weisz, a North Carolina State associate professor of crop science and small grains specialist; Kathy Flanders, an Auburn University associate professor of entomology; and Jeff Holland, a Purdue associate professor of entomology. Their findings were published in the *Journal of Economic Entomology*.

Provided by Purdue University

Citation: Wheat resistance genes failing, new approach needed to stop flies (2011, January 24) retrieved 24 April 2024 from <https://phys.org/news/2011-01-wheat-resistance-genes-approach-flies.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.