

## C60 could form a new kind of gel

February 17 2011, by Lisa Zyga



Simulation data of different structures: (a) a non-phase separated fluid, (b) a phase separated crystal, (c) a gel, and (d) isolated clusters. Image credit: C. Patrick Royall, et al.

(PhysOrg.com) --  $C_{60}$ , the spherical carbon molecule also known as a buckminsterfullerene, has intrigued scientists for its unique properties and potential applications in nanotechnology and electronics. Now



scientists have found that  $C_{60}$  may have another unusual property: it may take the form of a one-component gel under certain conditions. To date, all known gels consist of two components: an evenly distributed substance (a colloid) and a substance that dissolves the colloid (a solvent).

Scientists have previously discovered that  $C_{60}$  can take the form of different phases of matter, including solids and liquids. Here, chemists Patrick Royall from the University of Bristol and Stephen Williams from the Australian National University found that  $C_{60}$  can theoretically exist in a dense liquid phase containing clusters, which bind together to form a gel structure, specifically a "spinodal" gel. The gel is made entirely of carbon.

In their study, the scientists performed <u>computer simulations</u> showing that  $C_{60}$  can form a gel at moderately high temperatures and very high quench rates. The simulations showed that  $C_{60}$  gels form in about 10 nanoseconds and are stable at <u>room temperature</u> for at least 100 nanoseconds, which is the maximum time that the simulations were run. Although the gel showed some coarsening, the scientists predict that it would remain stable for more than 100 nanoseconds. Eventually, however, the gel would separate into a crystal and a gas.

As far as experimentally demonstrating the  $C_{60}$  gel, the scientists predict that it will be a challenge, largely due to the extremely high quench rates required, which are not currently experimentally feasible. However, they may investigate creative ways to lower the quench rate, or try to use larger fullerenes such as  $C_{540}$ , which may also become a carbon gel. In any case, the potential existence of a one-component <u>gel</u> could lead to an overall better understanding of the nature of gels.

**More information:** C. Patrick Royall and Stephen R. Williams. " $C_{60}$ : the first one-component gel?" <u>arXiv:1102.2959v1</u> [cond-mat.soft]



via: The Physics arXiv Blog

## © 2010 PhysOrg.com

Citation: C60 could form a new kind of gel (2011, February 17) retrieved 23 May 2024 from <u>https://phys.org/news/2011-02-c60-kind-gel.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.