

Fleeting fluctuations in superconductivity disappear close to transition temperature

February 13 2011



Ivan Bozovic

(PhysOrg.com) -- As part of an ongoing effort to uncover details of how high-temperature superconductors carry electrical current with no resistance, scientists at Johns Hopkins University and the U.S. Department of Energy's Brookhaven National Laboratory have measured fluctuations in superconductivity across a wide range of temperatures using terahertz spectroscopy. Their technique allows them to see fluctuations lasting mere billionths of a billionth of a second, and reveals that these fleeting fluctuations disappear 10-15 Kelvin (K) above the

transition temperature (T_c) at which superconductivity sets in.

"Our findings suggest that in cuprate superconductors, the transition to the non-superconducting state is driven by a loss of coherence among the [electron pairs](#)," said Brookhaven physicist Ivan Bozovic, a co-author on a paper describing the results in [Nature Physics](#) online, February 13, 2011.

Scientists have been searching for an explanation of high- T_c superconductivity in cuprates since these materials were discovered some 25 years ago. Because they can operate at temperatures much warmer than conventional superconductors, which must be cooled to near [absolute zero](#) (0 K or -273 degrees Celsius), high- T_c superconductors have the potential for real world applications. If scientists can unravel the current-carrying mechanism, they may even be able to discover or design versions that operate at room temperature for applications such as zero-loss power transmission lines. For this reason, many researchers believe that understanding how this transition to superconductivity occurs in cuprates is one of the most important open questions in physics today.

In conventional superconductors, electron pairs form at the transition temperature and condense into a collective, coherent state to carry current with no resistance. In high- T_c varieties, which can operate at temperatures as high as 165 K, there are some indications that electron pairs might form at temperatures 100-200 K higher, but only condense to become coherent when cooled to the transition temperature.

To explore the phase transition, the Johns Hopkins-BNL team sought evidence for superconducting fluctuations above T_c .

"These fluctuations are something like small islands or droplets of superconductivity, within which the electron pairs are coherent, which pop up here and there and live for a while and then evaporate to pop up

again elsewhere," Bozovic said. "Such fluctuations occur in every superconductor," he explained, "but in conventional ones only very, very close to T_c — the transition is in fact very sharp."

Some scientists have speculated that in cuprates, on the contrary, superconducting fluctuations might exist in an extremely broad region, all the way up to the temperature at which the electron pairs form. In the present study, the scientists tackle this question head-on, by measuring the conductivity as a function of temperature and frequency up to the terahertz range.

"With this technique, one can see superconducting fluctuations as short-lived as one billionth of one billionth of a second — the shortest possible — and over the entire phase diagram," Bozovic said.

The scientists studied a superconductor containing variable amounts of lanthanum and strontium layered with copper oxide. The samples were fabricated at Brookhaven, using a unique atomic-layer molecular beam epitaxy system that allows for digital synthesis of atomically smooth and perfect thin films. Terahertz spectroscopy measurements were performed at Johns Hopkins.

The central finding was somewhat surprising: The scientists clearly observed superconducting fluctuations, but these fluctuations faded out relatively quickly, within about 10-15 K above T_c , regardless of the lanthanum/strontium ratio.

This implies that in cuprates at the transition temperature, electron pairs lose their coherence. This is in contrast to what happens in conventional superconductors, where the [electron pairs](#) break apart at the [transition temperature](#).

"So, unlike in conventional superconductors, the transition in cuprates is

not driven by electron (de)pairing but rather by loss of coherence between pairs — that is, by phase [fluctuations](#)," Bozovic said. "The hope is that understanding this process in full detail may bring us one step closer towards cracking the enigma of high-temperature [superconductivity](#)."

More information: Temporal correlations of superconductivity above the transition temperature in $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ probed: [dx.doi.org/10.1038/nphys1912](https://doi.org/10.1038/nphys1912)

Provided by Brookhaven National Laboratory

Citation: Fleeting fluctuations in superconductivity disappear close to transition temperature (2011, February 13) retrieved 9 April 2024 from <https://phys.org/news/2011-02-fleeting-fluctuations-superconductivity-transition-temperature.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--