

Fruit flies can detect heavy hydrogen: study

February 16 2011, by Lin Edwards



In the present study wildtypes of different origin were shown to be quite different in their behavior towards odors of different origin. Credit: Rickard Ignell

(PhysOrg.com) -- A new study by researchers in Greece and the US has found that fruit flies can discriminate between normal and heavy hydrogen (deuterium) isotopes, which adds weight to a new theory of how the sense of smell works.

Dr Efthimios Skoulakis of the Alexander Fleming Biomedical Sciences Research Center in Vari, Greece and colleagues presented <u>fruit flies</u> (*Drosophila melanogaster*) with acetophenone, an odorant molecule often used in perfumes, in a T-shaped maze. In one branch the molecule contained normal hydrogen, while in the other the <u>hydrogen atoms</u> in the molecule were replaced by deuterium. The flies were able to choose the branch they preferred.



The fruit flies, which are known to have a strong sense of smell, showed a definite preference for the molecule with more hydrogen and their aversion to the deuterated molecule grew as more hydrogens were replaced. When fruit flies that had been genetically modified to have no sense of smell were used, they showed no preference.

The fruit flies' ability to discriminate between the molecules was also demonstrated by successfully giving the flies mild electric shocks to their feet as they walked on the floor of the maze to condition them to selectively avoid either form of the molecule.

Dr Skoulakis said the results support a new theory of olfaction first proposed in 1996 by co-author Dr Luca Turin, a biophysicist who is now at the Massachusetts Institute of Technology in Cambridge in the US. The new theory suggests odorants are detected by their vibrations rather than by their shape.

Deuterium differs from the normal isotope of hydrogen in that its nucleus contains a proton and a neutron instead of just a proton. This roughly doubles the mass of the deuterium atom, and so while it has similar properties to normal hydrogen, the larger mass means bonds between <u>deuterium</u> and other atoms within molecules vibrate more slowly.

The leading theory of how smell works is that odorant <u>molecules</u> are detected by receptor proteins in the olfactory membranes because of their shape, which fits into the shape of a cavity in the protein, like a key in a lock. If the molecule fits a signal is sent to the brain.

Dr Turin's theory is that the electrons of the odorant might be able to cross a receptor membrane only if the bonds in the molecule are vibrating at exactly the right frequency. Since an odorant with normal hydrogen will have different vibrational properties to the deuterated



odorant, the two odorants should smell different even though their shape is the same.

As further confirmation the researchers tested the fruit flies with nitriles, which have a similar vibrational frequencies to that of deuterium-carbon bonds. The fruit flies had an aversion to the nitriles as expected.

Humans have not been shown to have the ability to discriminate between compounds containing normal and heavy hydrogen, but Dr Turin said there were unpublished reports of a similar ability in at least one dog, which ignores an odorant it is trained to detect if the molecule is deuterated.

The paper is published in the journal *Proceedings of the National Academy of Sciences (PNAS)*.

More information: Molecular vibration-sensing component in Drosophila melanogaster olfaction, *PNAS*, Published online before print February 14, 2011, doi:10.1073/pnas.1012293108

© 2010 PhysOrg.com

Citation: Fruit flies can detect heavy hydrogen: study (2011, February 16) retrieved 19 April 2024 from https://phys.org/news/2011-02-fruit-flies-heavy-hydrogen.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.