

Giant ring of black holes

February 9 2011



X-ray: NASA/CXC/MIT/S.Rappaport et al, Optical: NASA/STScI

(PhysOrg.com) -- Just in time for Valentine's Day comes a new image of a ring -- not of jewels -- but of black holes.

This composite image of Arp 147, a pair of interacting galaxies located about 430 million light years from Earth, shows X-rays from the NASA's Chandra X-ray Observatory (pink) and optical data from the [Hubble Space Telescope](#) (red, green, blue) produced by the Space

Telescope Science Institute (STScI) in Baltimore, Md.

Arp 147 contains the remnant of a [spiral galaxy](#) (right) that collided with the [elliptical galaxy](#) on the left. This collision has produced an expanding wave of star formation that shows up as a blue ring containing in abundance of massive young stars. These stars race through their evolution in a few million years or less and explode as supernovas, leaving behind [neutron stars](#) and black holes.

A fraction of the neutron stars and black holes will have companion stars, and may become bright X-ray sources as they pull in matter from their companions. The nine X-ray sources scattered around the ring in Arp 147 are so bright that they must be [black holes](#), with masses that are likely ten to twenty times that of the Sun.

An X-ray source is also detected in the nucleus of the red galaxy on the left and may be powered by a poorly-fed supermassive black hole. This source is not obvious in the composite image but can easily be seen in the X-ray image. Other objects unrelated to Arp 147 are also visible: a foreground star in the lower left of the image and a background quasar as the pink source above and to the left of the red galaxy.

Infrared observations with NASA's [Spitzer Space Telescope](#) and ultraviolet observations with NASA's Galaxy Evolution Explorer (GALEX) have allowed estimates of the rate of star formation in the ring. These estimates, combined with the use of models for the evolution of binary stars have allowed the authors to conclude that the most intense star formation may have ended some 15 million years ago, in Earth's time frame.

These results were published in the October 1st, 2010 issue of *The Astrophysical Journal*. The authors were Saul Rappaport and Alan Levine from the Massachusetts Institute of Technology, David Pooley

from Eureka Scientific and Benjamin Steinhorn, also from MIT.

Provided by Chandra X-ray Center

Citation: Giant ring of black holes (2011, February 9) retrieved 23 April 2024 from <https://phys.org/news/2011-02-giant-black-holes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.