

Profiling malaria-causing parasites

February 7 2011

The majority of fatal cases of malaria are caused by infection with the parasite *Plasmodium falciparum*. Most at risk are young children and women who are pregnant. A team of researchers, led by Patrick Duffy, at the National Institutes of Health, Rockville, has now developed an approach to profile *P. falciparum* parasites in such a way that they are able to identify parasite genes associated with severe infection.

In the study, they identified a distinct subset of genes in <u>parasites</u> infecting <u>pregnant women</u>, including a gene whose protein product is already known to be important for severe disease in these at risk patients and is a well-established pregnancy malaria vaccine candidate.

They also identified a subset of genes that distinguished parasites infecting children from those infecting pregnant women. The team hopes that this information will provide new understanding of the nature of severe childhood malaria. Future experiments will attempt to validate the protein products of these <u>genes</u> as biomarkers of disease severity and targets for new intervention strategies.

More information: View this article at: www.jci.org/articles/view/4345 ... 7ba810e70af4526cbadd

Provided by Journal of Clinical Investigation

Citation: Profiling malaria-causing parasites (2011, February 7) retrieved 21 June 2024 from



https://phys.org/news/2011-02-profiling-malaria-causing-parasites.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.