

Residual dipolar couplings unveil structure of small molecules

February 17 2011

German chemists at the Technische Universitaet Muenchen and the Karlsruhe Institute of Technology introduced a new method for identifying chemical compounds. The approach they used is an improvement on nuclear magnetic resonance measurements -- for decades one of the most successful methods for determining the chemical structure of organic molecules. The results now published in the scientific journal *Angewandte Chemie* show a handy approach to structural data when classical methods of analysis fail.

The team of Professor Burkhard Luy from KIT and Junior Professor Stefan F. Kirsch from the TUM has now shown for the first time that certain NMR parameters, the so-called residual dipolar couplings (RDCs), can make a significant contribution towards determining the constitution of chemical compounds when traditional methods fail. To do this they embedded molecules of the compound in a gel which slightly constricts their mobility. By stretching the gel, the molecules can be aligned along a preferred orientation. While residual dipolar couplings average out in solution, they become measurable in such partially aligned samples and provide valuable structural information that can be used to build a model of the molecule.

To test this new approach to [chemical structure](#) determination the scientists examined a molecule whose atomic composition was known, but not the precise connectivities of the individual atoms in the molecule. The molecule was obtained using a unique reaction, so there were no precedents for its structure. Classical methods of analysis failed

because of the compactness of the molecule. In this particular case it was only possible to determine the structure by means of residual dipolar couplings, so that the newly acquired knowledge could be used to draw conclusions about the formation of the molecule – something that in the past could only be speculated about.

"This type of analysis will not be suitable for all structures in the future," said scientists Luy and Kirsch. "There will still be molecules whose structures will defy all attempts at unraveling, in spite of tremendous efforts and cutting-edge technologies. But this new method provides us with one further tool to help us unravel the structural mysteries of nature."

More information: Grit Kummerlöwe, Benedikt Crone; Manuel Kretschmer, Stefan F. Kirsch und Burkhard Luy: Residual Dipolar Couplings as a Powerful Tool for Constitutional Analysis: The Unexpected Formation of Tricyclic Compounds. *Angewandte Chemie International Edition*, scheduled to go online 17 or 18 February 2011. [DOI:10.1002/anie.2010007305](https://doi.org/10.1002/anie.2010007305)

Provided by Technische Universitaet Muenchen

Citation: Residual dipolar couplings unveil structure of small molecules (2011, February 17)
retrieved 25 April 2024 from
<https://phys.org/news/2011-02-residual-dipolar-couplings-unveil-small.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--