

Smithsonian scientists discover 7 new species of fish

February 4 2011



This is a male *Starksia langi* -- one of the seven new species of blenny discovered by Smithsonian scientist Carole Baldwin and her team. Credit: Smithsonian

Things are not always what they seem when it comes to fish -- something scientists at the Smithsonian Institution and the Ocean Science Foundation are finding out. Using modern genetic analysis, combined with traditional examination of morphology, the scientists discovered that what were once thought to be three species of blenny in the genus *Starksia* are actually 10 distinct species. The team's findings are published in the scientific journal *ZooKeys*, Feb. 3.

Starksia blennies, small (less than 2 inches) fish with elongated bodies, generally native to shallow to moderately deep rock and <u>coral reefs</u> in the western Atlantic and eastern Pacific oceans, have been well-studied for more than 100 years. It would have been reasonable to assume that there was little about the group left to discover. Modern <u>DNA barcoding</u> techniques, however, suggested otherwise. While trying to match larval stages of coral reef fish to adults through DNA, the team of scientists



noticed contradictions between the preliminary genetic data and the current species classification. Further investigation revealed that the team was dealing with many species new to science, including the new *Starksia* blennies.

"DNA analysis has offered science a great new resource to examine old questions," said Carole Baldwin, a zoologist at Smithsonian's National Museum of Natural History and lead author of the paper. "This discovery is a perfect example of how DNA barcoding is illuminating species that we've missed before, particularly small cryptic reef fishes like *Starksia* blennies. We don't know where we stand in terms of understanding species diversity, and our work suggests that current concepts may be surprisingly incomplete."



This is a female *Starksia robertsoni* -- one of the seven new species of blenny discovered by Smithsonian scientist Carole Baldwin and her team. Credit: Smithsonian

But DNA analysis cannot stand on its own—Baldwin and her team only recognize genetic lineages as species if they are supported by morphology. So traditional morphological analysis, such as comparing patterns of pigmentation and numbers of fin rays, is conducted to solidify their findings.

One interesting aspect of the research is that *Starksia* species that were thought to be broadly distributed throughout the Caribbean—as most



Caribbean reef fish species are—break up into multiple species with geographically restricted ranges. One species in the study, for example, was divided into three—a species in the east (Bahamas/Turks and Caicos), one in the south (Curacao, Netherlands Antilles) and another in the west (Belize, Central America). Baldwin predicts that other widespread species in the genus may also represent species complexes that break into multiple, geographically distinct species after further study. Furthermore, the team's DNA data suggest that other types of Caribbean fish (e.g., some gobies) may similarly represent species complexes comprising numerous new species, and traditional concepts of speciation in the Caribbean may need to be re-evaluated.



This is a male *Starksia williamsi* -- one of the seven new species of blenny discovered by Smithsonian scientist Carole Baldwin and her team. Credit: Smithsonian

The team's combined molecular and morphological approach has not only increased the number of currently recognized species, it serves as an example of the continuing nature of scientific discovery. Because the resiliency of marine populations to human exploitation may be linked to species richness, an improved understanding of the diversity and distribution of deep-reef life may be critical.

Provided by Smithsonian



Citation: Smithsonian scientists discover 7 new species of fish (2011, February 4) retrieved 23 April 2024 from https://phys.org/news/2011-02-smithsonian-scientists-species-fish.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.