

# Ancient 'hyperthermals' a guide to anticipated climate changes

March 16 2011

---



Sediment samples in the lab of Richard Norris obtained by the Ocean Drilling Program reveal the mark of "hyperthermals," warming events lasting thousands of years that changed the composition of the sediment and its color. The packaged sediment sample on the left contains sediment formed in the wake of a 55-million-year-old warming event and the sample on the right is sediment from a later era after global temperatures stabilized. Credit: Scripps Institution of Oceanography, UC San Diego

Bursts of intense global warming that have lasted tens of thousands of years have taken place more frequently throughout history than previously believe, according to evidence gathered by a team led by Scripps Institution of Oceanography, UC San Diego researchers.

Richard Norris, a professor of geology at Scripps who co-authored the report, said that releases of [carbon dioxide](#) sequestered in the deep oceans were the most likely trigger of these ancient "hyperthermal" events. Most of the events raised average [global temperatures](#) between 2°

and 3° Celsius (3.6 and 5.4° F), an amount comparable to current conservative estimates of how much temperatures are expected to rise in coming decades as a consequence of anthropogenic global warming. Most hyperthermals lasted about 40,000 years before temperatures returned to normal.

The study appears in the March 17 issue of the journal *Nature*.

"These hyperthermals seem not to have been rare events," Norris said, "hence there are lots of ancient examples of [global warming](#) on a scale broadly like the expected future warming. We can use these events to examine the impact of global change on marine ecosystems, climate and ocean circulation."

The hyperthermals took place roughly every 400,000 years during a warm period of Earth history that prevailed some 50 million years ago. The strongest of them coincided with an event known as the Paleocene-Eocene Thermal Maximum, the transition between two geologic epochs in which global temperatures rose between 4° and 7° C (7.2° and 12.6° F) and needed 200,000 years to return to historical norms. The events stopped taking place around 40 million years ago, when the planet entered a cooling phase. No warming events of the magnitude of these hyperthermals have been detected in the geological record since then.

Phil Sexton, a former student of Norris' now at the Open University in the United Kingdom, led the analysis of sediment cores collected off the South American coast. In the cores, evidence of the warm periods presented itself in bands of gray sediment layered within otherwise pale greenish mud. The gray sediment contained increased amounts of clay left after the calcareous shells of microscopic organisms were dissolved on the sea floor. These clay-rich intervals are consistent with ocean acidification episodes that would have been triggered by large-scale releases of carbon dioxide. Large influxes of carbon dioxide change the

chemistry of seawater by producing greater amounts of carbonic acid in the oceans.

The authors concluded that a release of carbon dioxide from the deep oceans was a more likely cause of the hyperthermals than other triggering events that have been hypothesized. The regularity of the hyperthermals and relatively warm ocean temperatures of the period makes them less likely to have been caused by events such as large melt-offs of methane hydrates, terrestrial burning of peat or even proposed cometary impacts. The hyperthermals could have been set in motion by a build-up of carbon dioxide in the deep oceans caused by slowing or stopping of circulation in ocean basins that prevented carbon dioxide release.

Norris noted that the hyperthermals provide historical perspective on what Earth will experience as it continues to warm from widespread use of fossil fuels, which has increased carbon dioxide concentrations in the atmosphere nearly 50 percent since the beginning of the Industrial Revolution. Hyperthermals can help scientists produce a range of estimates for how long it will take for temperatures to fully revert to historical norms depending on how much warming human activities cause.

"In 100 to 300 years, we could produce a signal on Earth that takes tens of thousands of years to equilibrate, judging from the geologic record," he said.

The scientists hope to better understand how fast the conditions that set off hyperthermals developed. Norris said that 50 million year old sediments in the North Sea are finely layered enough for scientists to distinguish decade-to-decade or even year-to-year changes.

Provided by University of California - San Diego

Citation: Ancient 'hyperthermals' a guide to anticipated climate changes (2011, March 16)  
retrieved 17 April 2024 from <https://phys.org/news/2011-03-ancient-hyperthermals-climate.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.