


# Contrast agent for tumor diagnostics: Phosphorescent metal-organic coordination polymers for optical imaging

March 23 2011

---



(PhysOrg.com) -- X-rays are not the only way: visible and especially infrared light can also be used to image human tissue. The effectiveness of optical imaging processes can be significantly improved with suitable dyes used as contrast agents. In the journal *Angewandte Chemie*, a team led by Wenbin Lin at the University of North Carolina has now introduced a novel contrast agent that marks tumor cells in vitro. The dye is a phosphorescent ruthenium complex incorporated into

nanoparticles of a metal–organic coordination polymer, which allows an extraordinarily high level of dye loading.

Fluorescent dyes accumulate in varying amounts in different types of tissue. Such contrast agents make it possible to use [optical imaging](#) to differentiate between healthy and tumorous tissue. However, this method is limited by the fact that very high concentrations of dye are needed to produce sufficiently strong fluorescence. Organic dye molecules packed at high concentrations into nanocapsules tend to quench each other's fluorescence. Materials that fluoresce more strongly, such as quantum dots, are often not biocompatible.

This team has now developed an alternative: metal complexes connected to form lattice-like coordination polymers. Coordination polymers are metal–organic structures consisting of metal ions, which act as connecting points, linked by bridges made of organic molecules or coordination complexes. The scientists made such polymers with bridges consisting of a light-emitting complex of the metal ruthenium. Zirconium ions proved to be suitable connecting points. These tiny structures form spherical nanoparticles.

The [ruthenium](#) complexes do not fluoresce, but rather phosphoresce, which means that they emit light for a proportional length of time after irradiation with light. Because they are not placed inside a nano-transport container, but are a component of the nanoparticle, it is possible to attain a very high level of dye loading—in this instance over 50 %. Quenching of the phosphorescence at high concentrations does not occur in such complexes.

In order to prevent the glowing particles from rapidly dissolving and to increase the biocompatibility, they were coated with thin layers of silicon dioxide and a layer of polyethylene glycol. The latter acts as an anchor point for anisamide, a molecule that specifically binds to

receptors that are far more common on the surfaces of many types of tumor cell than on healthy cells.

In a cell culture, it was possible to selectively mark a line of cancer cells with the phosphorescent [nanoparticles](#). The researchers hope that it will be possible to develop [contrast agents](#) for the use of optical imaging for tumor detection based on these new metal–organic nanomaterials.

**More information:** Wenbin Lin et al., Phosphorescent Nanoscale Coordination Polymers as Contrast Agents for Optical Imaging, *Angewandte Chemie International Edition*, [dx.doi.org/10.1002/anie.201008277](https://doi.org/10.1002/anie.201008277)

Provided by Wiley

Citation: Contrast agent for tumor diagnostics: Phosphorescent metal-organic coordination polymers for optical imaging (2011, March 23) retrieved 19 April 2024 from <https://phys.org/news/2011-03-contrast-agent-tumor-diagnostics-phosphorescent.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.