

Fossil bird study describes ripple effect of extinction in animal kingdom

March 7 2011, by Leeann Bright



Jessica Oswald, an NSF predoctoral fellow at the Florida Museum of Natural History on the UF campus, holds the mandible, or beak, of an extinct species of cowbird, *Pandanaris convexa*, recently discovered for the first time in Mexico. The bird has previously only been found at the Rancho La Brea fossil site in California and a site in Reddick, between Gainesville and Ocala in North Central Florida. Oswald is lead author of a new study in the March 8, 2011, print edition of the journal *Palaeogeography, Palaeoclimatology, Palaeocology*. The research shows the ripple effect throughout the animal kingdom caused by the extinction of large mammals 20,000 years ago, including the disappearance of a cowbird species. Image: Florida Museum of Natural History/UF photo by Kristen Grace

(PhysOrg.com) -- A University of Florida study demonstrates extinction's ripple effect through the animal kingdom, including how the demise of large mammals 20,000 years ago led to the disappearance of one species of cowbird.

The study shows the trickle-down effect the loss of large mammals has on other [species](#), and researchers say it is a lesson from the past that should be remembered when making conservation, game and land-use decisions today.

"There's nothing worse for a [terrestrial ecosystem](#) than the loss of large mammals – and the loss of apex predators like sharks, tuna and other large fish will have the same negative impact on the oceans," said study co-author David Steadman, ornithology curator at the Florida Museum of Natural History on the UF campus. "We're seeing it with the loss of lions and elephants in parts of Africa, as well as in Florida with the decline of panthers. There's no question these losses will have a negative domino effect on our ecosystems."

The fossil study of eight songbird species from northern Mexico by Florida Museum ornithologists is currently available online and will appear in Tuesday's print edition of the journal *Palaeogeography, Palaeoclimatology, Palaeocology*.

An extinct cowbird, *Pandanaris convexa*, is the most common bird found at the fossil site called T rapa, in Sonora, Mexico, about 150 miles south of Arizona. This is the first time fossils of the large bird, a member of the blackbird family, have been found in Mexico.

Finding the extinct cowbird at the fossil site was unpredictable and unexpected, according to Jim Mead, chair of the department of geosciences at East Tennessee State University, who has collected a variety of fossils at the site, including the birds used in the study. Mead

described the findings at T rapa as "bizarre and exciting."

"The tropical environment is unusual because the site is so far from the coast," Mead said. "The fossil record also provides evidence animals migrated from north to south and, unexpectedly, from south to north."

The cowbird has previously only been found at the Rancho La Brea fossil site in California and a site in Reddick, between Gainesville and Ocala in North Central Florida. The study expands the bird's known range and creates new questions about whether it may have lived across the southern U.S.

"The extinct cowbird needed grasslands and these big mammals to survive," said lead author Jessica Oswald, a National Science Foundation predoctoral fellow at the Florida Museum. "Those two things play into each other because mega mammals maintain grasslands. They keep big trees from coming in and colonizing the areas because they graze, stomp and trample little saplings."

Like modern cowbirds, this species probably fed on seeds and insects large mammals exposed, Oswald said. The mammals included extinct species of ground sloth, mammoth, horse, tapir, camel and bison.

About 20,000 years ago, most of these large mammals went extinct, which led to the extinction of scavengers like condors and vultures, as well as cowbirds, Steadman said. Extinctions, especially mass extinctions, can cause radical species loss and changes in species distribution.

"Big species can't exist in a vacuum, nor can smaller species," Steadman said. "When one piece of the puzzle goes extinct, there is no good way of predicting what sort of trickle-down effect, what kind of cascade effect that will have."

The study also confirms the area was once marshy grassland, possibly surrounded by a savanna near a river. Fossils of plants, reptiles and [mammals](#) of all sizes, and 31 species of birds other than songbirds have been recovered from the Térapa site over the past 10 years. Most of these species are found today in grasslands or wetlands, Steadman said.

Steadman and Oswald used the Florida Museum's more than 24,000 skeletal specimens of birds to identify the Mexican fossils.

Songbirds make up more than 50 percent of the world's living bird species, but the [fossil record](#) is poorly developed, especially in Central and South America. Oswald said this study helps build the [fossil](#) record of songbirds in Mexico.

Finding bird fossils, as well as bones of other small animals, is a time-consuming and labor-intensive process. Sediment is placed in a fine mesh sieve and water is used to remove dirt and debris from the bones.

Provided by University of Florida

Citation: Fossil bird study describes ripple effect of extinction in animal kingdom (2011, March 7) retrieved 19 April 2024 from

<https://phys.org/news/2011-03-fossil-bird-ripple-effect-extinction.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--