

# Back to the future with mummified trees

March 17 2011, By Lily Whiteman

---



The landscape of Quttinirpaaq National Park is dominated by glaciers and sparse high-arctic tundra vegetation. Credit: Joel Barker, Byrd Polar Research Center, Ohio State University

When in Quttinirpaaq National Park in the Canadian Arctic, Ohio State University Earth scientist Joel Barker initially spotted some pieces of dead trees scattered on the barren ground near a glacier. Immediately, he knew he had found something akin to a looking glass peering into the Arctic's ecological past.

The Hazen Plateau on Ellesmere Island, a polar desert where [winter temperatures](#) can currently dip down to 50 below zero, is currently too cold and dry to support forests; the only living trees that now dot the park's desolate landscape are dwarf willows. Therefore, the pieces of wood must have come from trees that lived millions of years ago, when the Arctic was still warm enough to support forests.

The trees probably died in a landslide, as indicated by deposits still present on surrounding material. The killing landslide toppled and engulfed the trees quickly enough to seal them from oxygen and to prevent water from circulating through. The result: The remaining pieces of dead trees are now in a perfectly preserved mummified state, with much of their [organic material](#) still intact. (By contrast, if the trees had been petrified, much of their organic material would have been slowly replaced by minerals and eventually turned to stone.)

In fact, the organic material in the dead trees is so well preserved that the wood can still burn, and even the most delicate tree structures, such as leaves, are present. "The dead trees look just like the dried-out dead wood lying outside now," said Barker.



The mummified trees have very narrow growth rings, indicating a very harsh environment during their lifetime. Credit: Joel Barker, Byrd Polar Research Center, Ohio State University

The summer following his discovery, Barker returned with colleagues, supported by an EAGER (EARly-concept Grants for Exploratory Research) grant from the National Science Foundation (NSF). Barker and Ohio State colleagues Yo Chin and David Elliot explored and

collected samples from the source of the scattered tree pieces--a deposit about 100 feet (30 meters) wide by 30 feet (10 meters) high.

Analyses of the organic material can reveal much about the source trees and about the ecological conditions that existed when they lived. For example, because pollen that disappeared about 12 million years ago are absent from the wood, Barker believes that the trees are probably at most 12 million years old. And because ocean cores that contain land-derived sediment indicate that forests disappeared from the Arctic about 2 million years ago, Barker believes that the wood is probably at least about that old.

Other evidence indicates to Barker that his wood samples came from trees that were once part of an ecologically stressed ecosystem. For example, the tree pieces have narrow growth rings, indicating that, when alive, the trees lacked sufficient nutrients for growth.

In addition, Barker said that only five different species of trees were represented in the ancient wood he collected. Such relatively low species diversity is typical of stressed ecosystems. That is because as an ecosystem becomes more stressed and less habitable, fewer species can survive in it. "Probably living at the most northern extent of its range, this forest was probably living at the edge, just hanging on, just barely staying alive," said Barker.

"The most stressed ecosystems are vulnerable to even small changes in climate," said Barker. "So studies of the sensitivity of stressed ecosystems to past cooling might improve our understanding of the sensitivity of ecosystems to stresses created by ongoing climate change." Such studies may, for example, help reveal the speed at which ecosystems will respond to current and future temperature changes.

Barker said that the mummified trees from Quttinirpaaq National

Park--which he found through a tip from a park ranger--may be more useful for climate change studies than any other mummified forests found in the Arctic so far. Why? Because all other mummified forests that have, so far, been found in the Canadian Arctic lived further south than the Quttinirpaaq trees. And a mummified forest that was found in Greenland--the only mummified forest found north of the Quttinirpaaq forest--had more species diversity than the Quttinirpaaq trees.

Therefore, these other mummified forests were probably not as stressed and were probably not as vulnerable to climate change as were the Quttinirpaaq trees. However, additional mummified forests will probably be exposed by retreating glaciers as [climate change](#) continues.

In the meantime, Barker will continue to analyze his mummified samples. He plans to extract individual components from pieces of mummified tree trunks to identify environmental conditions that the trees experienced on an annual basis throughout their lives. In addition, he plans to compare specific tree components, such as leaf cuticles, from mummified material with the same components from contemporary trees of the same species. Differences between the mummified and contemporary components of the same tree species may provide clues about the evolution of those species.

But with such experiments, Barker will only be literally scratching the surface of his mummified samples. He hopes to soon return to their site of origin to dig deeper to find their source forest, where trees may be standing upright in a so-called "growth position."

Provided by National Science Foundation

Citation: Back to the future with mummified trees (2011, March 17) retrieved 3 May 2024 from <https://phys.org/news/2011-03-future-mummified-trees.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.