


The rich chemistry around an evolved star

March 11 2011



A deep optical image of the carbon star IRC+10216, showing traces of its surrounding envelope. New SMA observations study the rich chemistry of the envelope, and find 442 spectral lines from over fifty molecules. Credit: Izan Leao; the Very Large Telescope

(PhysOrg.com) -- Over 170 molecules have been detected in space, from simple diatomic molecules like CO to complex organic molecules with over 70 atoms, like fullerene.

These [molecules](#) play a critical role in the development of [molecular clouds](#) as they form new stars and planetary systems, and of course in the chemistry that later develops on the surfaces of planets. One of the major issues in modern astronomy is figuring out exactly where all these molecules and associated [dust grains](#) came from.

The variable star CW Leo, also known as IRC+10216, is one of the brightest objects in the sky as seen from Earth; it is about 450 light-years away. It shines mostly in the infrared (not optical) because the central star is surrounded by a dense cloud of dust and gas that it ejected in a late stage of its evolution; that dust blocks the [optical light](#). The material is known to be rich in carbon-bearing molecules.

CfA astronomers Nimesh Patel, Ken Young, Carl Gottlieb, Pat Thaddeus, Bob Wilson, Mark Reid, Mike McCarthy, and Eric Keto, together with five colleagues, used the [Submillimeter Array](#) (SMA) to study the spectrum of IRC+10216 across a wavelength band, in an effort to detect and characterize as many molecules in the star's envelope as possible.

The scientists report finding an amazing 442 spectral lines in their survey, more than 200 of them detected for the first time in any astronomical source. All but 149 can be identified as arising from specific molecules. In addition to measuring the strengths of the lines and the motions of the molecules responsible, the SMA survey also obtained images of the nebula around in the star in the light of each of these species. The unidentified features, for example, tend to arise from compact regions around the star and probably correspond to hotter states of the known molecules; future work is needed to confirm this

conclusion. The new results provide a remarkable view of the rich chemistry around this nearby star, and help to strengthen the conclusion that many complex molecules trace their origin to the envelopes of evolved stars.

Provided by Harvard-Smithsonian Center for Astrophysics

Citation: The rich chemistry around an evolved star (2011, March 11) retrieved 2 May 2024 from <https://phys.org/news/2011-03-rich-chemistry-evolved-star.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
