

Agulhas leakage fueled by global warming could stabilize Atlantic overturning circulation: study

April 27 2011



This figure shows the Agulhas system and its leakage into the South Atlantic. The southward shift of the winds in a warming climate (right panel) cause a southward shift of the subtropical front (red arrows), enlarging the "gateway" for Agulhas leakage around the tip of Africa. Increased Agulhas leakage acts to enhance Atlantic overturning, which causes a feedback on climate. (Background colors are dynamic height integrated between 2000 m and the surface). Credit: Erik van Sebille, UM/RSMAS

(PhysOrg.com) -- The Agulhas Current which runs along the east coast of Africa may not be as well known as its counterpart in the Atlantic, the Gulf Stream. But now researchers are taking a closer look at this current and its "leakage" from the Indian Ocean into the Atlantic Ocean--and



what that may mean for climate change.

In results of a study published in this week's issue of the journal *Nature*, a team of scientists led by University of Miami Rosenstiel School of Marine & Atmospheric Science Oceanographer Lisa Beal, suggests that Agulhas leakage could be a significant player in global climate variability.

The Agulhas Current transports warm and salty waters from the tropical Indian Ocean to the southern tip of Africa. There most of the water loops around to remain in the Indian Ocean (the Agulhas Retroflection), while some water leaks into the fresher Atlantic Ocean via giant Agulhas rings.

Once in the Atlantic, the salty Agulhas leakage waters eventually flow into the Northern Hemisphere and act to strengthen the Atlantic overturning circulation by enhancing deep-water formation.

Atlantic overturning circulation is technically known as Atlantic meridional overturning circulation (AMOC); it carries warm shallow water into northern latitudes and returns cold deep water southward across the equator.

Recent research points to an increase in Agulhas leakage over the last few decades, caused primarily by human-induced climate change.

The finding is profound, oceanographers say, because it suggests that increased Agulhas leakage could trigger a strengthening in Atlantic overturning circulation--at a time when warming and accelerated meltwater input in the North Atlantic has been predicted to weaken it.

"This could mean that current IPCC model predictions for the next century are wrong, and there will be no cooling in the North Atlantic to



partially offset the effects of global climate change over North America and Europe," said Beal.



Current meters are secured to lab benches, ready for deployment into the Agulhas Current. Credit: Lisa Beal

"Instead, increasing Agulhas leakage could stabilize the oceanic heat transport carried by the Atlantic overturning circulation."

There are also paleoceanographic data to suggest that dramatic peaks in Agulhas leakage over the past 500,000 years may have triggered the end of glacial cycles.

These data are further evidence that the Agulhas system and its leakage play an important role in the planet's climate, Beal and others say.

"This study shows that local changes in atmospheric and oceanic conditions in the Southern Hemisphere can affect the strength of the ocean circulation in unexpected ways," said Eric Itsweire, director of the National Science Foundation (NSF)'s physical oceanography program, which funded the research.



"Under a warming climate," said Itsweire, "the Agulhas Current system near the tip of South Africa could bring more warm salty water from the Indian to the Atlantic Ocean and counteract opposing effects from the Arctic Ocean."

The study establishes the need for additional research in the region that focuses on Agulhas rings, as well as on the leakage, believes Beal.

Climate modeling experiments are critical, she said, and need to be supported by paleoceanographic data and sustained observations to firmly establish the role of the Agulhas system in a warming climate.



A current meter and chain of floats are deployed as part of several-kilometerslong mooring. Credit: Lisa Beal

"Our goal now is to get more of the scientific community involved in research on the Agulhas system and its global effects," said Beal. "The emphasis has been too long in the North Atlantic."

The Agulhas Current Time-Series Experiment, or ACT, was launched in April 2010 to measure the variability of the Agulhas Current using a combination of current meter moorings and satellite data.



Beal, who serves as chief scientist, spent one month aboard the research vessel Knorr in the southwest Indian Ocean deploying oceanographic instruments.

The data gathered in situ, when combined with along-track satellite information, will help increase our understanding of how the Agulhas system is changing in a warming climate, Beal said.

The scientific team included Beal, Wilhelmus P.M. de Ruijter of Utrecht University in the Netherlands, Arne Biastoch of Leibniz- Institut für Meereswissenschaften (IFM-GEOMAR) in Germany, and Rainer Zahn of the Universitat Autònoma de Barcelona in Spain.

It also included members of the Scientific Committee for Oceanic Research (SCOR) Working Group 136 on the Climatic Importance of the Agulhas System, sponsored by SCOR, the International Association for the Physical Sciences of the Oceans, and the World Climate Research Program.

Provided by National Science Foundation

Citation: Agulhas leakage fueled by global warming could stabilize Atlantic overturning circulation: study (2011, April 27) retrieved 3 May 2024 from https://phys.org/news/2011-04-agulhas-leakage-fueled-global-stabilize.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.