

# Boluochia closely related to longipteryx, study shows

April 6 2011

---



Figure 1. Photograph of the holotype of *Longipteryx chaoyangensis* (IVPP V12325). Scale bar represents 1 cm. Credit: O'Connor JK

Very little was known about Mesozoic birds until nearly complete skeletons began to be discovered in the now famous Jehol Group deposits of northeastern China.

The first few specimens to be found were partial skeletons, most preserving only the voids of the bones. Over the past two decades, new specimens have continued to be uncovered at an unprecedented rate. More recent published discoveries have typically been complete, including [feathers](#), and benefited from recent advances in preparational techniques, thus providing a huge wealth of data regarding species

diversity in northeastern [China](#) during the Early Cretaceous and the greatest wealth of data on Mesozoic birds anywhere in the world. Using this new data, a renewed look at the first few Jehol birds to be described reveals new information.

Paleornithologists of the Institute of [Vertebrate Paleontology](#) and [Paleoanthropology](#) (IVPP), Chinese Academy of Sciences, post-doctor Jingmai O'Connor, Director Zhou Zhonghe, and researcher Dr. Zhang Fucheng, reinvestigated *Boluochia zhengi*, one of the first birds described from the Jehol deposits. This taxon is very fragmentary, preserving clearly only the ankles and feet, however, this detailed study revealed that this species can now be assigned to the most diverse recognized group of enantiornithines, the Longipterygidae. Other members of this group (*Longipteryx*, *Longirostravis*, *Shanweinia* and *Rapaxavis*) are all known from nearly complete specimens and preserve elongate rostra with teeth restricted to the tips of their jaws. This is interpreted as a trophic specialization that would have made food items available to members of this clade that were inaccessible to other [birds](#), and may have contributed to their evolutionary success evident from their high [species diversity](#). Although only a few fragments of the skull are preserved in the only known specimen of *Boluochia*, O'Connor, Zhou and Zhang were able to identify subtle clues that hint at the presence of an elongate rostrum in this taxon as well. These, however, could not be recognized until more complete specimens were discovered and available for comparison.



Figure 2: Photographs of the holotype of *Boluochia zhengi* (IVPP V9770).  
Credit: O'Connor JK

There are two lineages of longipterygids: those that are smaller with reduced hands (*Longirostravis*, *Shanweinia* and *Rapaxavis*), and a more basal lineage with large claws on the hands and large teeth, before only represented by *Longipteryx*. Among longipterygids, O'Connor, Zhou and Zhang determined that the preserved morphology most closely aligns *Boluochia* with *Longipteryx*; unfortunately both taxa are from the younger Jiufotang Formation, and basal longipterygids (those with primitive morphologies such as claws on the hands such as *Longipteryx*) remain unknown from the older Yixian formation that has produced the apparently more derived *Longirostravis* and *Shanweinia*. *Boluochia* and *Longipteryx* share an unusual morphology of the foot, with metatarsals III and IV subequal in length, and a very large and robust pygostyle (the distal tail [bone](#)). However, *Boluochia* can still be distinguished from

Longipteryx and the two remain separate species based on the outwards deflection of the outermost foot bone (distal end of metatarsal IV).

Provided by Institute of Vertebrae Paleontology and Paleoanthropology

Citation: Boluochia closely related to longipteryx, study shows (2011, April 6) retrieved 14 August 2024 from <https://phys.org/news/2011-04-boluochia-longipteryx.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.