

# Chernobyl's radioactivity reduced the populations of birds of orange plumage

April 26 2011

---



Bird populations fell as the levels of radiation in peripheral zones of Chernobyl (Ukraine) rose. Credit: Rafael Palomo Santana

On April 26, 1986, history's greatest nuclear accident took place northwest of the Ukrainian city of Chernobyl. Despite the scale of the disaster, 25 years later, we still do not know its real effects. An international team of investigators has shown for the first time that the colour of birds' plumage may make them more vulnerable to radioactivity.

Radiation causes oxidative stress, damages biological molecules and may have "important" negative effects on organisms in relatively high doses, like those found in certain zones close to Chernobyl.

"In the case of the [birds](#) studied, these effects were seen in the size of their populations", says Ismael Galván, lead author of the study and researcher in the Laboratory of Ecology, Systematics and Evolution at the University of Paris-Sur, in France, speaking to SINC.

According to the study, which has been published in the journal *Oecologia*, bird populations fell as the levels of radiation in peripheral zones of Chernobyl (Ukraine) rose. In total, the researchers analysed the abundance of 97 bird [species](#) exposed to different levels of radiation during four years.

In the majority of the birds (64 species), the populations diminished with the level of radioactivity. "Nevertheless, the populations of a few species (the 33 remaining species) experienced positive effects from the radiation (though the magnitude of these effects was very low in some cases), perhaps due to the reduction in competition with other species", explains Galván.

Colour: a bird's weak or strong point

The scientists concentrated on the colouring generated by melanins – pigments which protect from ultraviolet radiation and generate camouflage patterns – of the nearly one hundred species of bird studied. The reason: the type of pigmentation may interfere with the ability to resist radioactivity's negative effects.

"The impact on the populations depends, at least in part, on the amount of plumage whose colouring is generated by pheomelanin, one of the two main types of melanins, which produces orangish and brownish colours", the Spanish expert adds.

The birds of [Chernobyl](#) with the most pheomelanism (with the most plumage coloured by pheomelanin) were judged to be the "most

negatively" affected by the radioactivity. As the pigment consumes glutathione (one of the antioxidants most susceptible to radiation and whose level tends to be diminished by its effects), in these birds, the capacity to combat the oxidative stress generated by [radiation](#) "probably" diminishes.

**More information:** Galván, Ismael; Mousseau, Timothy A.; Moller, Anders P. "Bird population declines due to radiation exposure at Chernobyl are stronger in species with pheomelanin-based coloration" *Oecologia* 165(4): 827-835, April 2011.

Provided by FECYT - Spanish Foundation for Science and Technology

Citation: Chernobyl's radioactivity reduced the populations of birds of orange plumage (2011, April 26) retrieved 27 April 2024 from <https://phys.org/news/2011-04-chernobyl-radioactivity-populations-birds-orange.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--