


New test for germs: Fluorescing DNazymes detect metabolic products from bacteria

April 7 2011



(PhysOrg.com) -- Germs in food, bioterrorism, drug-resistant bacteria and viruses—these are the problems of our time that make early detection of pathogens particularly important. Whereas conventional methods are either slow or require complex instruments, Yingfu Li and a team at McMaster University in Hamilton (Ontario, Canada), additionally supported by the Sentinel Bioactive Paper Network, have now developed an especially simple, universal fluorescence test system that specifically and rapidly detects germs by means of their metabolic products. As the researchers report in the journal *Angewandte Chemie*, It isn't even necessary to know which substance the test is reacting to.

Traditionally [germs](#) have been detected through microbiological methods, which are very precise but can take days or weeks. PCR- or antibody-based methods are rapid but require many steps and special equipment. “We were motivated to develop an especially simple, but

very rapid and precise method,” says Li. “It must also be universal, meaning that it should be possible to develop tests for any desired germ using the same principle.”

“When a pathogen is metabolically active and multiplying in a given medium, it releases many substances into this environment. These are what we want to use,” says Li. The idea is to produce DNAszymes that react to a pathogen-specific product. A DNAszyme is a synthetic one-stranded DNA molecule with catalytic activity. Making a large pool of DNA molecules with random sequences and subjecting these to repeated selection and amplification steps allows for the development of molecules with the desired property.

At the core of the conceptual DNAszyme is a single RNA nucleotide. To its right and left are a fluorescing dye and a quencher. A quencher is a molecule that switches off the fluorescence of a dye when it is nearby. The researchers developed a DNAszyme that binds to a specific metabolic product from *E. coli* bacteria, which causes the DNAszyme to change its shape. In this altered form, the DNAszyme has RNA-splitting capability and cuts its own strand at the location of the RNA nucleotide. This separates the quencher from the dye, which begins to fluoresce. The fluorescence indicates that *E. coli* is present in the sample. This DNAszyme does not react to other bacteria.

“Through targeted selection, it should be possible to find a specific DNAszyme for any desired germ,” says Li. “It is not necessary to know what the metabolic product is, or to isolate it from the sample.” By using a common cell culture step, it is possible for the pathogens in a sample to multiply before the test, which allows for detection of a single original cell.

More information: Yingfu Li, Fluorogenic DNAszyme Probes as Bacterial Indicators, *Angewandte Chemie International Edition* 2011, 50,

No. 16, 3751–3754, [dx.doi.org/10.1002/anie.201100477](https://doi.org/10.1002/anie.201100477)

Provided by Wiley

Citation: New test for germs: Fluorescing DNAzymes detect metabolic products from bacteria (2011, April 7) retrieved 8 April 2024 from <https://phys.org/news/2011-04-germs-fluorescing-dnazymes-metabolic-products.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.