


Researchers use GPS data to model effects of tidal loads on Earth's surface

April 15 2011, By Katie Neith



The outline of each ellipse represents the motion the ground makes as the earth flexes in response to the time and space dependent ocean tides. There is an ellipse for each of the GPS sites used in the study, and the color indicates the amplitude of the tidal response movement.

(PhysOrg.com) -- For many people, Global Positioning System (GPS) satellite technology is little more than a high-tech version of a traditional paper map. Used in automobile navigation systems and smart phones,

GPS helps folks find their way around a new neighborhood or locate a nearby restaurant. But GPS is doing much, much more for researchers at the California Institute of Technology (Caltech): it's helping them find their way to a more complete understanding of Earth's interior structure.

Up until now, the best way to explore Earth's internal structures—to measure geological properties such as density and elasticity—has been through seismology and laboratory experiments. "At its most fundamental level, seismology is sensitive to specific combinations of these properties, which control the speed of seismic waves," says Mark Simons, professor of geophysics at Caltech's Seismological Laboratory, part of the Division of Geological and Planetary Sciences. "However, it is difficult using seismology alone to separate the effects that variations in density have from those associated with variations in elastic properties."

Now Simons and Takeo Ito, visiting associate at the Seismological Laboratory and assistant professor of earth and planetary dynamics at Nagoya University in Japan, are using data from GPS satellite systems in an entirely new way: to measure the solid earth's response to the movements of ocean tides—which place a large stress on Earth's surface—and to estimate separately the effects of Earth's density and the properties controlling response when a force is applied to it (known as elastic moduli).

Their work was published in this week's issue of *Science Express*.

By using measurements of Earth's movement taken from high-precision, continuously recording permanent GPS receivers installed across the western United States by the Plate Boundary Observatory (PBO), the researchers were able to observe tide-induced displacements—or movements of Earth's surface—of as little as one millimeter. PBO is a component of EarthScope, a program that seeks to understand the

processes controlling earthquakes and volcanoes by exploring the structure and evolution of the North American continent.

The team focused on understanding the properties of the asthenosphere, a layer of weak and viscous upper mantle that lies below Earth's crust, and used those measurements to build one-dimensional models of Earth's response to the diurnal tides in the western United States.

"The asthenosphere plays an important role in plate tectonics, as it lies directly under the plates," explains Ito. "The results of our study give us a better understanding of the asthenosphere, which in turn can help us understand how the plates move."

Still shot from a movie that shows the variation in the sea surface associated with the one constituent of the ocean tide, a pattern that repeats about every 12 hours and 25.2 minutes. This variation in sea surface height generates a periodic load on the earth, which is what was measured in the study. To watch the full movie, [click here](#).

The models provided a look at the variations in density from Earth's surface down to a depth of about 400 kilometers. The researchers found that the density of the asthenosphere under the western United States and the eastern Pacific Ocean is abnormally low relative to the global average.

"Variations in density can either be caused by variations in the chemical makeup of the material, the presence of melt, or due to the effects of thermal expansion, whereby a given material will decrease in density as its temperature increases," explains Simons. "In this study, we interpret the observed density anomaly to be due to the effects of elevated temperatures in the asthenosphere below the western United States and neighboring offshore areas. The required peak temperature anomaly would be about 300 degrees Celsius higher than the global average at

those depths."

This type of data provides keys to understanding the chemical and mechanical dynamics of the planet, such as how heat flows through the mantle and how tectonic plates on Earth's surface are evolving.

"It is amazing that by measuring the twice-a-day centimeter-scale cyclic movement of Earth's surface with a GPS receiver, we can infer the variation of density 220 kilometers below the surface," says Simons.

Now that the researchers know it is possible to use GPS to derive measurements of internal [Earth](#) structures, they anticipate several new directions for this research.

"We hope to extend the observations to be global in scope, which may require temporary deployments of GPS in important areas that are typically tectonically bland—in other words, devoid of significant earthquakes and volcanoes—and thus do not have existing dense continuous GPS arrays already in place," says Simons. Next steps may also include going beyond the current one-dimensional depth-dependent models to build 3-D models, and combining the GPS approach with more conventional seismic approaches.

"The method we developed for gathering data from [GPS](#) devices has significant potential for improving 3-D images of Earth's internal structure," says Ito.

The findings of the study, "Probing asthenospheric density, temperature and elastic moduli below the Western United States," are available online in Science Express and will be featured in a future issue of the journal *Science*.

More information: The findings of the study, "Probing asthenospheric

density, temperature and elastic moduli below the Western United States," are available online in *Science Express* and will be featured in a future issue of the journal *Science*.

Provided by California Institute of Technology

Citation: Researchers use GPS data to model effects of tidal loads on Earth's surface (2011, April 15) retrieved 9 April 2024 from <https://phys.org/news/2011-04-gps-effects-tidal-earth-surface.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--