


Scientists discover mechanism that could feed solar explosions

April 15 2011



(a) Top row: SOT Ca H-line images showing the dark prominence bubble developing a sharp protrusion that develops into the large plume. The plume is fully separated from the bubble by 15:38:43 UT. The outlines of the bubble and plume are traced in these images. (b) Middle row: SDO/AIA 17.1 nm images of the bubble and plume development. The yellow dashed lines are taken from the Hinode/SOT tracings in the top row. They show that the bubble and plume brightness in this spectral line correspond exactly to the dark bubble and plume outlines in the SOT images. (c) Bottom row: SDO/AIA 19.3 nm images of the bubble and plume development. The black dashed lines are the outlines taken from the top row. Again, the brightness of the bubble and plume in this spectral line are apparent.

(PhysOrg.com) -- Coronal Mass Ejections (CMEs) are violent solar explosions that can propel up to 10 billion tons of the Sun's atmosphere – at a million miles an hour – out through the corona and into space. These fast, powerful ejections can take as little as 18 hours to reach Earth and give rise to geomagnetic storms, which can disrupt radio transmissions, induce large currents in power lines and oil pipelines, seriously disrupt spacecraft and be extremely hazardous to astronauts. New instruments on advanced spacecraft have provided fresh insight into these cataclysmic phenomena, and illuminated a path toward predicting space weather.

In a paper published today in the journal *Nature*, researchers from the Solar and Astrophysics Laboratory (LMSAL) of the Lockheed Martin Advanced Technology Center (ATC), along with colleagues at the Harvard-Smithsonian Center for Astrophysics, Kyoto University, and the High Altitude Observatory of the National Center for Atmospheric Research (which is sponsored by the National Science Foundation) – have discovered a turbulent convective flow system in solar "quiescent" prominences suspended in the corona—the Sun's outer atmosphere– that point to a mechanism by which hot coronal plasma (and presumably magnetic flux) are injected upwards into the coronal cavity system. Coronal cavities are large magnetic flux ropes suspended in the corona, typically in the polar regions of the Sun. These flux ropes all eventually erupt in the form of CMEs that can impact the interplanetary and terrestrial space environments.


“How these large flux ropes erupt is a poorly understood fundamental process in the science of space weather. Our discovery points to a way in which intermittent ‘bubbles’ in solar prominences can inject new mass and magnetic flux into the flux ropes, thus slowly building up their magnetic buoyancy over time. These ‘bubbles’, which can be as wide as

several Earth diameters, are analogous to the blobs of material in a Lava Lamp that are heated by a light from below, become buoyant, and rise to the top to deposit their energy, then drop back down again. By this mechanism coronal cavity flux ropes could grow slowly until they are able to exceed the ‘tethering’ forces of overlying magnetic fields and thus erupt as CMEs,” said Dr. Thomas Berger, lead author of the Nature paper, and solar physicist at the Lockheed Martin Solar and Astrophysics Lab at the ATC. “If we can show in further research that the prominence bubbles are indeed magnetic flux emergence events taking place below prominences, we can verify that we’ve found a new mechanism for transferring magnetic flux from the convection zone into the corona, and perhaps establish a predictive tool for the eruption of CMEs based on the rate of observed flux injection.”

The researchers used observations from the Atmospheric Imaging Assembly (AIA) on NASA’s recently launched Solar Dynamics Observatory (SDO) and NASA’s Focal Plane Package for the Solar Optical Telescope (SOT) on the Japanese Hinode satellite. Both instruments were designed and built at the ATC.

It was the high spatial and temporal resolution of SOT, combined with the broad temperature coverage of AIA that unlocked the mystery. SOT movies from 2006--2009 reveal dark "bubbles" forming below 10,000 K prominence material. These bubbles go unstable and form turbulent upflow plumes that rise into the prominence and the 1,000,000 K "coronal cavity" above the prominence. SOT images alone couldn't identify the source of the bubbles' buoyancy—was it magnetic field concentration or thermal energy that led to the buoyancy relative to the heavy prominence above? In August 2010, using a simultaneous prominence observation by SOT and the newly launched AIA instrument suite, Berger and his team discovered that the bubbles were heated to temperatures of at least 250,000 K and more likely 1,000,000 K before rising into the prominence. This is 25-100 times hotter than the

overlying prominence and implies that in addition to any magnetic buoyancy in the system, there is significant thermal buoyancy as well.



(a) Hinode X-Ray Telescope (XRT) image of the region surrounding the prominence. The coronal cavity above the prominence is denoted by the dashed black lines. XRT images coronal plasma in the 1--3 MK temperature range. Note the much larger field-of-view of XRT compared to the following images. The axes are labelled in arcseconds from solar disk center. (b) Hinode Solar Optical Telescope (SOT) image of the quiescent prominence below the coronal cavity. The image is taken in the spectral line of ionized Calcium at 396.8 nm in the visible spectrum (the "H-line"). This spectral line images plasma in the 7,000--10,000 K temperature range. The white box shows the field-of-view of Figure 2 below. The dark prominence bubble and a large plume from the bubble are pointed out in the image. Note that the image has been rotated relative to (a) to place the solar limb approximately horizontal. (c) Solar Dynamics Observatory (SDO) Atmospheric Imaging Assembly (AIA) image of the same prominence as in (b). The image is taken in the spectral line of ionized Helium at 30.4 nm. This spectral line images plasma in the 50,000--80,000 K temperature range. The white dashed line defines a line along which emission is measured in Figure 3. (d) SDO/AIA image of the prominence in the spectral line of ionized iron at 17.1 nm showing the prominence plasma at a temperature of approximately 700,000 K. Note that the prominence bubble and plume are bright in this

spectral line. (e) SDO/AIA image of the prominence in the spectral line of ionized iron at 19.3 nm showing the prominence plasma at a temperature of approximately 1,250,000 K. Note that the prominence bubble and plume are bright in this spectral line. (f) SDO/AIA image of the prominence in the spectral line of ionized iron at 21.1 nm showing the prominence plasma at a temperature of approximately 1,700,000 K. Note that the prominence bubble and plume are bright in this spectral line. Axes in panels (b)–(f) are labelled in Megameters (1,000,000 meters). The UT times of the images are shown in the upper right corners.

“This discovery is significant because it revises the common view that the magnetic field in the corona dominates the gas pressure and allows only simple, laminar, flows along magnetic field lines. Here we establish that the prominence bubbles and resultant plumes are a form of the Rayleigh-Taylor instability, a buoyant turbulent flow system that, combined with the cool downflowing plasma in quiescent prominences, represents a form of convection, or overturning motion in the prominence/corona system – the first confirmed discovery of convection in the solar outer atmosphere,” added Berger. “It is apparent that our understanding of basic forces at work in the corona must be revised to include turbulent motions that can deform the magnetic field lines and produce novel flow and mixing systems.”

The Solar and Astrophysics Laboratory at the ATC conducts basic research into understanding and predicting space weather and the behavior of our [Sun](#) including its impacts on Earth and climate. It has a 47-year-long heritage of spaceborne solar instruments including the Soft X-ray Telescope on the Japanese Yohkoh satellite, the Michelson Doppler Imager on the ESA/NASA Solar and Heliospheric Observatory, the solar telescope on NASA’s Transition Region and Coronal Explorer, the Focal Plane Package on the Japanese Hinode satellite, the Solar X-ray Imagers on GOES-N, -O and -P, the Extreme Ultraviolet Imager

instruments on NASA's twin STEREO spacecraft, and the Helioseismic and Magnetic Imager and the Atmospheric Imaging Assembly on NASA's Solar Dynamics Observatory. The ATC is currently building both the science instrument and spacecraft for NASA's Interface Region Imaging Spectrometer (IRIS), a Small Explorer Mission scheduled for launch in late 2012.

The ATC is the research and development organization of Lockheed Martin Space Systems Company (LMSSC).

Provided by Lockheed Martin Corporation

Citation: Scientists discover mechanism that could feed solar explosions (2011, April 15)
retrieved 28 April 2024 from

<https://phys.org/news/2011-04-scientists-mechanism-solar-explosions.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--