


Ocean acidification will likely reduce diversity, resiliency in coral reef ecosystems: new study

May 29 2011



A new study of Papua New Guinea's "champagne reefs" in *Nature Climate Change* by the University of Miami, the Australian Institute of Marine Science and the Max-Planck Institute for Marine Microbiology in Germany concludes that ocean acidification, along with increased ocean temperatures, will likely severely reduce the diversity and resilience of coral reef ecosystems within this century. These reefs provide sobering illustrations of how coral reefs may look in 100 years if ocean acidification conditions continue to worsen. Credit: Katharina Fabricius/Australian Institute of Marine Science

A new study from University of Miami (UM) Rosenstiel School of Marine & Atmospheric Science scientists Chris Langdon, Remy Okazaki and Nancy Muehllehner and colleagues from the Australian Institute of Marine Science and the Max-Planck Institute for Marine Microbiology

in Germany concludes that ocean acidification, along with increased ocean temperatures, will likely severely reduce the diversity and resilience of coral reef ecosystems within this century.

The research team studied three natural volcanic CO₂ seeps in Papua New Guinea to better understand how [ocean acidification](#) will impact coral reefs ecosystem diversity. The study details the effects of long-term exposure to high levels of carbon dioxide and low pH on Indo-Pacific coral reefs, a condition that is projected to occur by the end of the century as increased man-made CO₂ emissions alter the current pH level of seawater, turning the oceans acidic.

"These 'champagne reefs' are natural analogs of how [coral reefs](#) may look in 100 years if ocean acidification conditions continue to get worse," said Langdon, UM Rosenstiel School professor and co-principal investigator of the study.

The study shows shifts in the composition of coral species and reductions in biodiversity and recruitment on the reef as pH declined from 8.1 to 7.8. The team also reports that reef development would cease at a pH below 7.7. The IPCC 4th Assessment Report estimates that by the end of the century, ocean pH will decline from the current level of 8.1 to 7.8, due to rising atmospheric CO₂ concentrations.

"The seeps are probably the closest we can come to simulating the effect of man-made CO₂ emissions on a coral reef," said Langdon. "They allow us to see the end result of the complex interactions between species under acidic ocean conditions."

The reefs detailed in this study have healthy reefs nearby to supply larvae to replenish the reefs. If pH was low throughout the region -- as projected for year 2100 -- then there would not be any healthy reefs to reseed damaged ones, according to Langdon.

More information: "Losers and winners in coral reefs acclimatized to elevated carbon dioxide concentrations," *Nature Climate Change*, June 2011.

Provided by University of Miami

Citation: Ocean acidification will likely reduce diversity, resiliency in coral reef ecosystems: new study (2011, May 29) retrieved 9 April 2024 from <https://phys.org/news/2011-05-ocean-acidification-diversity-resiliency-coral.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--