

Study: Rare deep-sea starfish stuck in juvenile body plan

May 2 2011



Scientists leveraged computer systems at the Ohio Supercomputer Center to help support their contention that class-level status of *Xyloplax* does not reflect their evolutionary history. This female specimen was collected along the Juan de Fuca Ridge off the coast of the state of Washington; it measures 4 mm in diameter (less than .25 inch) and shows brooded embryos. Credit: Ben Grupe/Scripps Institution of Oceanography, University of California, San Diego

A team of scientists has combined embryological observations, genetic sequencing, and supercomputing to determine that a group of small disk-shaped animals that were once thought to represent a new class of animals are actually starfish that have lost the large star-shaped, adult body from their life cycle.



In a paper for the journal *Systematic Biology* (sysbio.oxfordjournals.org), Daniel Janies, Ph.D., a computational biologist in the department of <u>Biomedical Informatics</u> at The Ohio State University (OSU), leveraged computer systems at the Ohio Supercomputer Center (OSC) to help support his contention that class-level status of *Xyloplax* does not reflect their <u>evolutionary history</u>.

"Although *Xyloplax* does not represent a new class, it an even more interesting animal now because it represents a rare example of how <u>natural selection</u> can shape the whole the life cycle," he explained. "By omitting the large adult stage, *Xylopax* found how to make a living in the nooks and crannies of sunken timbers on the deep-sea floor."

Janies collaborated on the paper with co-authors Janet R. Voight, Ph.D., in the department of Zoology at the Field Museum of Natural History in Chicago, and Marymegan Daly, Ph.D., in the department of Evolution, Ecology and Organismal Biology at Ohio State.

Janies and his colleagues are examining echinoderms – <u>starfish</u>, sea urchins and their close relatives – as part of a study for the United States' National Science Foundation's Assembling the Tree of Life project. Several OSU scientists, including Janies and Daly, have won these grants, typically valued at \$3 million over five years, to better understand the interrelationships of all forms of life.





A team of scientists has determined that small disk-shaped animals called *Xyloplax* are actually starfish that have lost the large star-shaped, adult body from their life cycle. Specimens shown in this image measure between 2 and 5 millimeters (less than .25 inch) in diameter, and were collected in 2010 from the Pacific Ocean at a depth of 2,202 meters (7,267 feet). Credit: Ben Grupe/Scripps Institution of Oceanography, University of California, San Diego

As tree-of-life analyses are computationally difficult, the current project is enabled by the computational muscle of OSC's flagship system, the 9,500-node IBM 1350 Opteron "Glenn Cluster."

"Last year, OSC deployed a \$4 million expansion to the Glenn Cluster with a specific focus on supporting researchers, like Drs. Janies and Daly, in Ohio's growing biosciences sector," said Ashok Krishnamurthy, interim co-executive director of OSC. "The center provides a worldclass computing environment for amazing bioscience research projects at places like OSU's Department of Biomedical Informatics, the Research Institute at Nationwide Children's Hospital and the Cleveland Clinic."



The researchers examined both genetic and anatomical data to support their hypothesis, comparing *Xyloplax* with a total of 86 species representing major lineages of the five living classes of echinoderms (sea cucumbers, sea urchins, starfish, brittle stars, and sea lilies). For accurate comparisons, they applied several analytical methods to search for the shared mutations and changes in anatomy among the lineages.

The results describe the relationships in a phylogenetic tree representing the best-supported hypothesis on how *Xyloplax* evolved from other echinoderms. Phylogenetics is the study of the evolutionary relationships and changes among various biological species as they evolve from a common ancestor.

Prior to this study, molecular investigations of *Xyloplax* were limited, because specimens in two early collections in the 1980s were fixed in a solution which degrades DNA. Voight recently collected specimens of *Xyloplax* in the northeastern Pacific Ocean and carefully preserved them in ethanol. From these specimens, Daly was able to sequence several genes for *Xyloplax*. The newer specimens of *Xyloplax* also included several brooding females containing embryos, providing Janies with an unprecedented view of the creature's early development.

Janies and colleagues theorize that *Xyloplax* differs from other starfish species because it is progenetic – that is, it has a rare, truncated <u>life cycle</u> that leaves the mature organism with features retained from its juvenile stages. For example, the arms of a starfish typically grow axially, like spokes of a wheel, as they develop from juveniles to adults, whereas *Xyloplax* grows along its circumference, like the wheel itself, and never develops arms.

"Irrespective of method or data sampling scheme, our results show that *Xyloplax* evolved from within starfish," Janies concluded. "*Xyloplax* is just a little starfish that has a strange body plan and habitat, so strange



that many could not recognize it as a starfish until we unlocked its genome and development."

Provided by Ohio Supercomputer Center

Citation: Study: Rare deep-sea starfish stuck in juvenile body plan (2011, May 2) retrieved 1 May 2024 from <u>https://phys.org/news/2011-05-rare-deep-sea-starfish-stuck-juvenile.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.