

Scientists track evolution and spread of deadly fungus, one of the world's major killers

May 3 2011

New research has shed light on the origins of a fungal infection which is one of the major causes of death from AIDS-related illnesses. The study, published today in the journal *PLoS Pathogens*, funded by the Wellcome Trust and the BBSRC, shows how the more virulent forms of *Cryptococcus neoformans* evolved and spread out of Africa and into Asia.

[Cryptococcus neoformans](#) is a species of often highly aggressive fungi. One particular strain of the [fungus](#) – known as *Cryptococcus neoformans* variety *grubii* (*Cng*) – causes meningitis amongst patients with compromised immune systems following HIV infection. There are believed to be over up to a million cases of cryptococcal meningitis each year, resulting in over 600,000 deaths. Infection with the fungus, which invades the central nervous system, is treated with a life long therapy of antifungal drugs, which can have highly unpleasant side effects.

Sitali Simwami and Dr Matthew Fisher from Imperial College London, together with colleagues from St Georges, University of London, Naresuan University, Thailand, and the CBS Fungal Biodiversity Centre, The Netherlands, used genetic sequencing techniques to compare the genetic diversity of *Cng* in 183 samples taken from the clinic and the environment in Thailand against the 77 samples from a global database. Thailand has an emerging HIV epidemic and nearly one in five HIV-infected patients are affected by cryptococcal infection.

"Cryptococcal meningitis kills hundreds of thousands of people each year, almost as many as malaria, yet gets little attention," explains Dr Fisher. "We know very little about where it originated from and how it evolved. If we can track its evolution and diversity, then we can begin to understand where the pathogen originates from, how it infects people and how it adapts to become more – or less – virulent. This information will be valuable in helping us identify potential therapeutic targets in the future."

The researchers found that *Cng* in Thailand exhibits significantly less genetic diversity in comparison to other areas of the world, especially Africa where many different lineages of the pathogen occur. This suggests that populations of the fungus in Africa will have a wider spectrum of virulent strains and higher rates of adaptation to antifungal treatments, implying that clinicians need to pay particular attention to the risk of drug-resistant forms of the fungus here.

Their analysis also suggested that the pathogen was introduced from Africa to Asia at some point within the last 7,000 years. Many human infectious diseases are thought to have emerged within the last 11,000 years, following the rise of agriculture and domestication of animals. In particular, it supports the idea that the pathogen was imported via infected pigeons, which were domesticated around 5,000 years ago. The common pigeon, which originated in Africa, is considered to be a carrier and potential spreader of the fungus, its faeces being a common environmental source of *Cng*.

Provided by Wellcome Trust

Citation: Scientists track evolution and spread of deadly fungus, one of the world's major killers (2011, May 3) retrieved 2 May 2024 from <https://phys.org/news/2011-05-scientists-track-evolution-deadly-fungus.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.