


Heavy metal meets hard rock: Battling through the ocean crust's hardest rocks

June 30 2011



A granoblastic basalt viewed under the microscope (picture is 2.3 mm across). Magnification shows a rock formed of small rounded mineral grains annealed together (plagioclase: white, pyroxene: light green and light brown, and magnetite or ilmenite: black). They may look inoffensive, but these rocks are the hardest material ever drilled in more than four decades of scientific ocean drilling. The rocks are very abrasive and aggressive to the drilling and coring tools, and difficult to penetrate. However, the samples recovered provide a treasure trove of information, recording the rocks' initial crystallization as a basaltic dike then their reheating at the top of the mid-ocean ridge magma chamber. These rocks represent the heat exchanger where thermal energy from the cooling and solidifying melt in the magma chamber below is exchanged with seawater infiltrating from the oceans, leading to the "black smoker-type" hot (>350°C) water vents on the seafloor. Credit: IODP/USIO

Integrated Ocean Drilling Program (IODP) Expedition 335 Superfast

Spreading Rate Crust 4 recently completed operations in Ocean Drilling Program (ODP) Hole 1256D, a deep scientific borehole that extends more than 1500 meters below the seafloor into the Pacific Ocean's igneous crust – rocks that formed through the cooling and crystallization of magma, and form the basement of the ocean floor.

An international team of scientists led by co-chief scientists Damon Teagle (National Oceanographic Center Southampton, University of Southampton in the UK) and Benoît Ildefonse (CNRS, Université Montpellier 2 in France) returned to ODP Hole 1256D aboard the scientific research vessel, JOIDES Resolution, to sample a complete section of intact oceanic [crust](#) down into gabbros.

This expedition was the fourth in a series and builds on the efforts of three expeditions in 2002 and 2005.

Gabbros are coarse-grained intrusive rocks formed by the slow cooling of basaltic magmas. They make up the lower two-thirds of the ocean crust. The intrusion of gabbros at the mid-ocean ridges is the largest igneous process active on our planet with more than 12 cubic kilometers of new magma from the mantle intruded into the crust each year. The minerals, chemistry, and textures of gabbroic rocks preserve records of the processes that occur deep within the Earth's mid-ocean ridges, where new ocean crust is formed.

"The formation of new crust is the first step in Earth's plate tectonic cycle," explained Teagle. "This is the principal mechanism by which heat and material rise from within the Earth to the surface of the planet. And it's the motion and interactions of Earth's tectonic plates that drive the formation of mountains and volcanoes, the initiation of earthquakes, and the exchange of elements (such as carbon) between the Earth's interior, oceans, and atmosphere."



Granoblastic dike samples were recovered in abundance by fishing tools during successive hole remediation operations. Science party members Sumiyo Miyashita and Yoshiko Adashi from Niigata University, Japan, examine large rock samples from Hole 1256D. Credit: IODP/USIO

"Understanding the mechanisms that construct new tectonic plates has been a major, long-standing goal of scientific ocean drilling," added Ildefonse, "but progress has been inhibited by a dearth of appropriate samples because deep drilling (at depths greater than 1000 meters into the crust) in the rugged lavas and intrusive rocks of the ocean crust continues to pose significant technical challenges."

ODP Hole 1256D lies in the eastern equatorial [Pacific Ocean](#) about 900 kilometers to the west of Costa Rica and 1150 kilometers east of the present day East Pacific Rise. This hole is in 15 million year old crust that formed during an episode of "superfast" spreading at the ancient East Pacific Rise, when the newly formed plates were moving apart by more than 200 millimeters per year (mm/yr).

"Although a spreading rate of 200 mm/yr is significantly faster than the

fastest spreading rates on our planet today, superfast-spread crust was an attractive target," stated Teagle, "because seismic experiments at active mid-ocean ridges indicated that gabbroic rocks should occur at much shallower depths than in crust formed at slower spreading rates. In 2005, we recovered gabbroic rocks at their predicted depth of approximately 1400 meters below the [seafloor](#), vindicating the overall 'Superfast' strategy."

Previous expeditions to Hole 1256D successfully drilled through the erupted lavas and thin (approximately one-meter-wide) intrusive "dikes" of the upper crust, reaching into the gabbroic rocks of the lower crust. The drilling efforts of Expedition 335 were focused just below the 1500-meter mark in the critical transition zone from dikes to gabbros, where magma at 1200°C exchanges heat with super-heated seawater circulating within cracks in the upper crust. This heat exchange occurs across a narrow thermal boundary that is perhaps only a few tens of meters thick.

In this zone, the intrusion of magma causes profound textural changes to the surrounding rocks, a process known as contact metamorphism. In the mid-ocean ridge environment this results in the formation of very fine-grained granular rocks, called granoblastic basalts, whose constituent minerals recrystallize at a microscopic scale and become welded together by magmatic heat. The resulting metamorphic rock is as hard as any formation encountered by ocean drilling and sometimes even tougher than the most resilient of hard formation drilling and coring bits.



Laying out a fishing tool on the rig floor. An illustration of the hard work by the drill ship crew Credit: IODP/USIO, Photo by Benoît Ildefonse

Expedition 335 reentered Hole 1256D more than five years after the last expedition to this site. The expedition encountered and overcame a series of significant engineering challenges, each of which was unique, although difficulties were not unexpected when drilling in a deep, uncased, marine [borehole](#) into igneous rocks.

The patient, persistent efforts of the drilling crew successfully cleared a major obstruction at a depth of 920 that had initially prevented reentry into the hole to its full depth of 1507 meters. Then at the bottom of the hole the very hard granular rocks that had proved challenging during the previous Superfast expedition were once more encountered. Although there may only be a few tens of meters of these particularly tenacious granoblastic basalts, their extreme toughness once more proved challenging to sample—resulting in the grinding down of one of the hardest formation coring bits into a smooth stump.

A progressive, logical course of action was then undertaken to clear the bottom of the hole of metal debris from the failed coring bit and drilling cuttings. This effort required the innovative use of hole-clearing

equipment such as large magnets, and involved over 240 kilometers of drilling pipe deployments (trips) down into the hole and back onto the ship. (The total amount of pipe "tripped" was roughly equivalent to the distance from Paris to the English coast, or from New York City to Philadelphia, or Tokyo to Niigata). These efforts returned hundreds of kilograms of rocks and drill cuttings, including large blocks (up to 5 kilograms) of the culprit granoblastic basalts that hitherto had only been very poorly recovered through coring. A limited number of gabbro boulders were also recovered, indicating that scientists are tantalizingly close to breaking through into the gabbroic layer.

Expedition 335 operations also succeeded in clearing Hole 1256D of drill cuttings, much of which appear to have been circulating in the hole since earlier expeditions.

"We recovered a remarkable sample suite of granoblastic basalts along with minor gabbros, providing a detailed picture of a rarely sampled, yet critical interval of the oceanic crust," Ildefonse observed. "Most importantly," he added, "the hole has been stabilized and cleared to its full depth, and is ready for deepening in the near future."

More information: [iodp.tamu.edu/scienceops/exped...
fast_rate_crust.html](http://iodp.tamu.edu/scienceops/exped...fast_rate_crust.html)

Provided by Integrated Ocean Drilling Program Management
International

Citation: Heavy metal meets hard rock: Battling through the ocean crust's hardest rocks (2011, June 30) retrieved 26 April 2024 from <https://phys.org/news/2011-06-heavy-metal-hard-ocean-crust.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.