


# New imaging tech promising for diagnosing cardiovascular disease, diabetes

June 9 2011, by Emil Venere



Researchers have developed a new type of imaging technology to diagnose cardiovascular disease and other disorders by measuring ultrasound signals from chemical bonds in molecules exposed to a pulsing laser. This "vibrational photoacoustic" image shows plaque in an arterial wall. Credit: Purdue University Weldon School of Biomedical Engineering image/Han-Wei Wang and Ji-Xin Cheng

Researchers have developed a new type of imaging technology to diagnose cardiovascular disease and other disorders by measuring ultrasound signals from molecules exposed to a fast-pulsing laser.

The new method could be used to take precise [three-dimensional images](#) of plaques lining arteries, said Ji-Xin Cheng, an associate professor of biomedical engineering and chemistry at Purdue University.

Other imaging methods that provide molecular information are unable to

penetrate tissue deep enough to reveal the three-dimensional structure of the plaques, but being able to do so would make better diagnoses possible, he said.

"You would have to cut a cross section of an artery to really see the three-dimensional structure of the plaque," Cheng said. "Obviously, that can't be used for living patients."

The imaging reveals the presence of carbon-hydrogen bonds making up [lipid molecules](#) in arterial plaques that cause heart disease. The method also might be used to detect fat molecules in muscles to diagnose diabetes and for other lipid-related disorders, including neurological conditions and [brain trauma](#). The technique also reveals nitrogen-hydrogen bonds making up proteins, meaning the imaging tool also might be useful for diagnosing other diseases and to study collagen's role in scar formation.

"Being able to key on specific chemical bonds is expected to open a completely new direction for the field," Cheng said

Findings are detailed in a paper to be published in [Physical Review Letters](#) and expected to appear in the June 17 issue. The findings represent the culmination of four years of research led by Cheng and doctoral student Han-Wei Wang.

The new technique uses nanosecond [laser pulses](#) in the near-infrared range of the spectrum. The laser generates molecular "overtone" vibrations, or wavelengths that are not absorbed by the blood. The pulsed laser causes tissue to heat and expand locally, generating pressure waves at the ultrasound frequency that can be picked up with a device called a transducer.

"We are working to miniaturize the system so that we can build an

endoscope to put into blood vessels using a catheter," Cheng said. "This would enable us to see the exact nature of plaque formation in the walls of arteries to better quantify and diagnose cardiovascular disease."

Lihong Wang, Gene K. Beare Distinguished Professor of Biomedical Engineering at Washington University in St. Louis, is a pioneer of using the "photoacoustic" imaging of blood vessels based on the absorption of light by the electrons in hemoglobin.

The Purdue researchers are the first to show that a strong photoacoustic signal can arise from the absorption of light by the [chemical bonds](#) in molecules. The near-infrared laser causes enough heating to generate ultrasound but not enough to damage tissue.

"You can measure the time delay between the laser and the ultrasound waves, and this gives you a precise distance, which enables you to image layers of the tissues for three-dimensional pictures," Cheng said. "You do one scan and get all the cross sections. Our initial target is cardiovascular disease, but there are other potential applications, including diabetes and [neurological conditions](#)."

The approach represents a major improvement over another imaging technique, called coherent anti-Stokes Raman scattering, or CARS, which has been used by the Purdue-based lab to study three-dimensional plaque formation in arteries.

Also leading the research are Michael Sturek, chair of the Department of Cellular and Integrative Physiology at the Indiana University School of Medicine; Robert P. Lucht, Purdue's Ralph and Bettye Bailey Professor of Combustion in Mechanical Engineering; and David Umulis, a Purdue assistant professor of agricultural and biological engineering. Other authors of the paper include Purdue graduate students Ning Chai, Pu Wang and Wei Dou and Washington University postdoctoral researcher

Song Hu.

Findings are based on research with pig tissues in laboratory samples and also with live fruit flies.

"You can see fat inside fly larvae, representing the potential to study how obesity affects physiology in humans," Cheng said.

**More information:** [Label-free Bond-selective Imaging by Listening to Vibrationally Excited Molecules](#)

## ABSTRACT

We report the realization of vibrational photoacoustic (VPA) microscopy using optical excitation of molecular overtone vibration and acoustic detection of the resultant pressure transients. Our approach eliminates the tissue scattering problem encountered in near-infrared spectroscopy and enables depth-resolved signal collection. The second overtone of the CH bond stretch around 8300  $\text{cm}^{-1}$ , where blood interference is minimal, is excited. We demonstrate 3-D VPA imaging of lipid-rich atherosclerotic plaques by excitation from the artery lumen, and lipid storage in live *Drosophila* larvae, with mm-scale penetration depth.

Provided by Purdue University

Citation: New imaging tech promising for diagnosing cardiovascular disease, diabetes (2011, June 9) retrieved 10 April 2024 from <https://phys.org/news/2011-06-imaging-tech-cardiovascular-disease-diabetes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---