

Methane gas from cows -- the proof is in the poo

June 6 2011

Scientists could have a revolutionary new way of measuring how much of the potent greenhouse gas methane is produced by cows and other ruminants, thanks to a surprising discovery in their poo.

Researchers from the University of Bristol and the Teagasc Animal and Grassland Research Centre in Ireland, have found a link between methane production and levels of a compound called archaeol in the faeces of several fore-gut fermenting animals including <u>cows</u>, sheep and deer.

The compound could potentially be developed as a biomarker to estimate the methane production from domestic and <u>wild animals</u>, allowing scientists to more accurately assess the contribution that ruminants make to global <u>greenhouse gas emissions</u>.

Co-author Dr Fiona Gill, who conducted the work as a postdoctoral researcher at Bristol and is now at the University of Leeds, said: "When it comes to calculating carbon budgets there is currently a lot of uncertainty surrounding animal methane contributions, particularly from wild ruminants.

"We're quite good at measuring man-made CO2 emissions, but techniques to measure the animal production of methane – a much more <u>potent greenhouse gas</u> – have serious limitations.

"If we can identify a simple <u>biomarker</u> for methane production in animal



stools, then we can use this along with information on diet and animal population numbers to estimate their total contribution to global methane levels."

Cows, sheep and other ruminants are thought to be responsible for around one-fifth of global methane production but the precise amount has proved difficult to quantify. Methane production from animals is often measured using respiration chambers, which can be laborious and are unsuitable for grazing animals.

Archaeol is thought to come from organisms called archaea, which are symbiotic or 'friendly' microbes that live in the foregut of ruminant animals. These microbes produce methane as a by-product of their metabolism and this is then released by the animal as burping and flatulence.

Principal investigator, Dr Ian Bull of Bristol's School of Chemistry said: "We initially detected archaeol in the faeces of several foregut fermenters including camels, cows, giraffes, <u>sheep</u> and llamas. We then expanded the study to evaluate the quantities of this compound in the faeces of cows with different diets.

"Two groups of cows were fed on different diets and then their <u>methane</u> <u>production</u> and faecal archaeol concentration were measured. The animals that were allowed to graze on as much silage as they wanted emitted significantly more methane and produced <u>faeces</u> with higher concentrations of archaeol than those given a fixed amount of silage, supplemented by concentrate feed.

"This confirms that manipulating the diet of domestic livestock could also be an important way of controlling methane gas emissions."

More information: The research is published in the journal Animal



Feed Science and Technology.

Provided by University of Leeds

Citation: Methane gas from cows -- the proof is in the poo (2011, June 6) retrieved 4 May 2024 from <u>https://phys.org/news/2011-06-methane-gas-cows-proof.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.