

Seeing the wood for the trees: New study shows sheep in tree-ring records

July 27 2011



These are sheep in the Norwegian mountains. Credit: Atle Mysterud

Nibbling by herbivores can have a greater impact on the width of tree rings than climate, new research has found. The study, published this week in the British Ecological Society's journal *Functional Ecology*, could help increase the accuracy of the tree ring record as a way of estimating past climatic conditions.

Many factors in addition to climate are known to affect the tree ring record, including attack from [parasites](#) and herbivores, but determining how important these other factors have been in the past is difficult.

Working high in the mountains of southern Norway, midway between Oslo and Bergen, a team from Norway and Scotland fenced off a large area of mountainside and divided it into different sections into each of

which a set density of domestic sheep was released every summer.

After nine summers, cross sections of 206 birch trees were taken and tree ring widths were measured. Comparing these with local temperature and the numbers of sheep at the location where the tree was growing allowed the team to disentangle the relationship between temperature and browsing by sheep and the width of [tree rings](#).

According to lead author Dr James Speed of the NTNU Museum of Natural History and [Archaeology](#): "We found tree ring widths were more affected by sheep than the [ambient temperature](#) at the site, although temperatures were still visible in the tree ring records. This shows that the density of herbivores affects the tree ring record, at least in places with slow-growing trees."

The impact of large herbivores on tree rings has, until now, been largely unknown, so these findings could help increase the accuracy of the tree ring record as a way of estimating past [climatic conditions](#), says Dr Speed: "Our study highlights that other factors interact with climate to affect tree rings, and that to increase the accuracy of the tree ring record to estimate past climatic conditions, you need to take into account the history of wild and domestic herbivores. The good news is that past densities of herbivores can be estimated from historic records, and from the fossilised remains of spores from fungi that live on dung."

"This study does not mean that using tree rings to infer past climate is flawed as we can still see the effect of temperatures on the rings, and in lowland regions tree rings are less likely to have been affected by [herbivores](#) because they can grow out of reach faster," he explains.

Tree rings give us a window into the past, and have been widely used as climate recorders since the early 1900s. The growth rings are visible in tree trunk cross sections, and are formed in seasonal environments as the

wood is laid down faster in summer than winter. In years with better growing conditions (in cool locations this usually means warmer) tree rings are wider, and because trees can be very long-lived and wood is easily preserved, for example in bogs and lakes, this allows very long time-series to be established, and climatic conditions to be estimated from the ring widths.

More information: James D. M. Speed, Gunnar Austrheim, Alison J. Hester and Atle Myrnes (2011), 'Browsing interacts with climate to determine tree ring increment', [doi: 10.1111/j.1365-2435.2011.01877.x](https://doi.org/10.1111/j.1365-2435.2011.01877.x), is published in *Functional Ecology* on 27 July 2011.

Provided by Wiley

Citation: Seeing the wood for the trees: New study shows sheep in tree-ring records (2011, July 27) retrieved 24 June 2024 from <https://phys.org/news/2011-07-wood-trees-sheep-tree-ring.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.