


Bragg reflectivity of X-rays: At the limit of the possible

August 23 2011



Absolute reflectivity of 13.9-keV x-rays from the (008) atomic planes of a diamond crystal in Bragg backscattering (a)-(c), and x-ray Lang transmission topogram of the diamond crystal (d). The diamond crystal, nearly defect-free in a volume of $5 \times 5 \times 1 \text{ mm}^3$, reflects more than 99% of hard x-ray photons backwards in Bragg diffraction, with a remarkably small variation in magnitude across the sample.

(PhysOrg.com) -- Researchers utilizing high-brightness x-rays at the U.S. Department of Energy Office of Science's Advanced Photon Source at Argonne National Laboratory have demonstrated that synthetic, nearly

defect-free diamond crystals can reflect more than 99% of hard x-ray photons backward in Bragg diffraction, with a remarkably small variation in the magnitude of reflectivity across the sample. Their results, which are published in *Nature Photonics*, represent a quantum leap to the largest reflectivity measured, at the limit of what is theoretically possible. It was attained under the most challenging conditions of normal incidence and with extremely high-energy x-ray photons.

Ultra-high-reflectance mirrors are essential components of the most sophisticated optical instruments, which are designed to cover the entire frequency spectrum and find use in a wide range of applications, from telescopes for astronomy to high-energy x-ray facilities such as the APS. In the x-ray regime, super-polished mirrors with close to 100% [reflectivity](#) are routinely used at grazing angles of incidence. However, at large angles of incidence, and especially at normal incidence, such high reflectivity had not been achieved.

The research in this study was conducted by a team comprising investigators from Argonne National Laboratory and the Technological Institute for Superhard and Novel Carbon Materials (TISNCM) in Russia. This investigation was motivated by the need for high-reflectivity x-ray mirrors at close-to-normal incidence for use in x-ray free electron laser oscillators (XFELs).

XFELs are x-ray light sources of the future that will provide scientists with fully coherent [x-rays](#) with record high spectral purity and average brightness, qualities that are highly desirable for experiments. Unlike high-gain x-ray lasers, XFELs are designed to be low-gain machines, requiring a low-loss optical cavity with high-reflectivity x-ray mirrors close to backscattering, similar to conventional table-top lasers.

The feasibility of XFELs was questioned by many experts because of

the high-reflectivity x-ray mirror limitations. These limitations are now removed by the demonstration of very high (more than 99%) reflectivity of hard x-rays in Bragg backscattering from diamond crystals.

Theoretical analysis presented in the Nature Photonics paper establishes that diamond has the highest (higher than any other crystal) Bragg reflectivity due to the uniquely small ratio of the extinction length in Bragg diffraction to the absorption length. Even in backscattering, where the reflectivity is lowest, it was thought to be more than 99%, and this has now been demonstrated.

Synthetic diamond crystals of sufficient size and crystalline perfection for use as high-reflectivity x-ray mirrors can be grown using modern techniques. In this experiment, synthetic diamond single crystals of very high crystal quality were grown at the TISNCM via the temperature gradient method under high-pressure and high-temperature conditions. The reflectivity measurements were performed at the X-ray Science Division beamlines 30-ID and 7-ID at the APS utilizing 23.7-keV and 13.9-keV photons, respectively.

Systematic observations of the more than 99% reflectivity show that high-reflectivity mirror limitations in the regime of hard x-rays have been eliminated. This is an important step toward realization of x-ray cavities for XFELs. Also, the results in this study support a broad range of potential applications of diamond crystals for high-reflectance, coherence-preserving, and resilient x-ray optics, such as x-ray monochromators, beam splitters, delay lines, high-finesse Fabry-Perot resonators, etc.

More information: Kwang-Je Kim, Yuri Shvyd'ko, and Sven Reiche “A Proposal for an X-Ray Free-Electron Laser Oscillator with an Energy-Recovery Linac”, *Phys. Rev. Lett.* 100, 244802 (2008).

Yuri Shvyd'ko, et al., “Near-100% Bragg reflectivity of X-rays”, *Nature*

Photonics, (published online August 2011).

[DOI:10.1038/nphoton.2011.197](https://doi.org/10.1038/nphoton.2011.197)

Provided by Argonne National Laboratory

Citation: Bragg reflectivity of X-rays: At the limit of the possible (2011, August 23) retrieved 25 April 2024 from <https://phys.org/news/2011-08-bragg-x-rays-limit.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.