

INTEGRAL observations suggest unified model for Active Galactic Nuclei requires a rethink

August 3 2011



Artist's impression of an AGN according to the unified model. Credit: ESA/NASA, the AVO project and Paolo Padovani

Scrutinizing a large sample of Active Galactic Nuclei with INTEGRAL, astronomers have found that, unexpectedly, sources affected by stronger absorption at lower energies show an excess emission in the hard X-ray band when compared to their less obscured counterparts. The excess is likely due to reflection of X-rays off the same dense clouds responsible for absorption. Unaccounted for in the unified paradigm of Active Galactic Nuclei, the finding calls for a rethinking of some of this model's facets. The reflected radiation could also represent the long-sought missing part of the Cosmic X-ray Background.

Supermassive [black holes](#) lie at the core of all [massive galaxies](#), including the [Milky Way](#). Whereas the majority of them are inactive, a small fraction of these black holes are accreting the surrounding matter at extraordinary rates. This causes them to radiate profusely across the entire electromagnetic spectrum. Known as [Active Galactic Nuclei](#) (AGN), these sources are often so bright that they outshine their [host galaxies](#) and have been detected out to the far reaches of the observable Universe.

Astronomers explain the wide variety of features discerned in different classes of AGN in terms of the anisotropic geometry of the black hole's immediate surroundings. The accreting black hole is fed via a disc, and a thick, obscuring torus of gas and dust is believed to encompass the disc and to absorb part of the radiation it emits.

Depending on the orientation of the torus with respect to an observer's line of sight, the view to the [galactic nucleus](#) may be obstructed to varying degrees. This 'unified' model, which provides a phenomenological interpretation of the observed AGN diversity, is now questioned, at least in part, by a new study performed with ESA's INTEGRAL observatory.

"The unified model predicts that all AGN exhibit the same behaviour when observed at hard X-ray wavelengths, regardless of the different emission they might show in other bands," explains Claudio Ricci, a PhD student from the Data Centre for Astrophysics (ISDC) at the University of Geneva, Switzerland. Ricci is the lead author on the paper reporting on the study and published online on 2 August 2011 in *Astronomy & Astrophysics*. "Absorption by the torus is weaker for photons with increasing energy, and it should have no effect at all, for the objects in our sample, on the hard X-ray photons probed by INTEGRAL," he adds.

To test this framework, Ricci and colleagues put together a sample of

165 AGN observed with INTEGRAL during its first 8 years of operation at energies between 20 and 250 keV. The sample comprises AGN subject to absorption at lower energies and also unabsorbed ones. "We expected to see no difference between classes, but instead we noticed two very distinct trends in the data," he notes.

Surprisingly, the sources affected by stronger absorption at lower energies (i.e. from the infrared to the soft X-ray bands) show an excess of emission, with respect to their less obscured counterparts, in the energy range 30–60 keV. No appreciable difference is observed at higher energies. The excess emission is a signature of X-rays being reflected off neutral hydrogen gas in the dense clouds that surround the black hole and disc, possibly the same clouds responsible for absorption at lower energies.

"A significantly stronger reflection in a certain class of AGN indicates that the environment of these objects is different," comments co-author Roland Walter, who is the Principal Investigator of the INTEGRAL team at the ISDC. "The discrepancy in the high-energy emission properties of different AGN types is unaccounted for in the leading theoretical framework and calls for a rethinking of some of its details," he adds.

In the light of the new data, the unified model's basic assumption that the same central engine powers all AGN is safe. However, the presumed existence, in all of them, of an anisotropic, toroidal structure needs revision. "We argue that various classes of AGN are characterised by a substantially different distribution of absorbing material around the black hole," notes Walter.

The team advocate a patchy but overall isotropic morphology for the absorber. In this scenario, the density of clouds around the central source would determine the amount of reflected X-rays, with denser

environments resulting in stronger reflection. The astronomers speculate about whether such morphological variations could characterise objects at different evolutionary stages, although the issue is still unclear and the subject of further investigation.

The discovery of X-ray reflection by AGN has another, far-reaching implication in the debate about the Cosmic X-ray Background (CXB), a tenuous, diffuse background radiation that pervades the X-ray sky from all directions. The CXB consists of the cumulative X-ray radiation emitted by all unresolved active galaxies across cosmic history. As this diffuse background peaks in the hard X-ray regime probed by INTEGRAL, identifying its individual sources has been among the mission's main science goals ever since it began operations.

"One of the observatory's milestones was the first accurate measurement of the CXB intensity in the energy range where its emission is stronger," notes Chris Winkler, INTEGRAL Project Scientist at ESA. Achieved in 2006, this result did not solve, but in fact deepened, a long-standing debate about the origin of the CXB. The measured intensity appeared to be far greater than that which astronomers could predict by modelling observations of individual AGN and summing their contributions. To match the measured value, a much larger number of heavily obscured AGN than has been observed was required.

Now, the study by Ricci and his colleagues proposes a fresh solution to the puzzle: when the reflected X-rays are added to the total budget of radiation sources in the CXB, invoking sources that have never been observed may no longer be necessary.

"The result relies on several years of data acquired with the most sensitive hard X-ray telescope currently in operation," comments Winkler. "By showing us where the missing part of the CXB lies, INTEGRAL might have solved a 30-year long mystery," he concludes.

More information: C. Ricci, et al., "Reflection in Seyfert Galaxies and the Unified Model of AGN", *Astronomy & Astrophysics*, 2011, 532, A102

Provided by European Space Agency

Citation: INTEGRAL observations suggest unified model for Active Galactic Nuclei requires a rethink (2011, August 3) retrieved 19 April 2024 from <https://phys.org/news/2011-08-galactic-nuclei-requires-rethink.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.