

## Researchers discover oldest evidence of nails in modern primates

## August 15 2011

From hot pink to traditional French and Lady Gaga's sophisticated designs, manicured nails have become the grammar of fashion.

But they are not just pretty — when nails appeared on all fingers and toes in modern primates about 55 million years ago, they led to the development of critical functions, including finger pads that allow for sensitive touch and the ability to grasp, whether it's a nail polish brush or remover to prepare for the next trend.

In a new study co-authored by University of Florida scientists, researchers recovered and analyzed the oldest fossil evidence of fingernails in modern primates, confirming the idea nails developed with small body size and disproving previous theories nails evolved with an increase in primate body size. More than 25 new specimens of Teilhardina brandti – an extinct primate originally described from a single lower molar — include pieces of upper teeth and ankle bones that show the mammal lived in trees. Its nails allowed the lemur-like animal to grasp onto branches and move through the trees with more agility, researchers said.

"If you take all the primates that are alive today, they're all going to have characteristics that look the same, but unlike people, many of them live in trees," said co-author Jonathan Bloch, an associate curator of vertebrate paleontology at the Florida Museum of Natural History on the UF campus. "By finding parts of the skeleton of this primitive primate, we are able to test whether nails were present in the common ancestor of



the group that includes lemurs, monkeys, and humans — it's direct evidence as opposed to speculation."

Appearing in the current online edition of the American Journal of Physical Anthropology, the study provides a better understanding of the evolutionary relationships of one of the oldest known modern primates, as well as the time frame and environmental conditions that allowed for the development of nails on all fingers and toes, an exclusive feature among primates.

Specimens of *T. brandti* were collected over the last seven years in northwestern Wyoming's Bighorn Basin and represent the earliest North American species from the group of euprimates, also known as "true" primates. The fossils date to the early Eocene epoch, about 55.8 million years ago, at the same time as a 200,000-year global warming event known as the Paleocene-Eocene Thermal Maximum occurred, Bloch said. Mammals evolved to be smaller during that time, when even- and odd-toed hoofed mammals, distantly related to modern deer and horses, also first appeared in the fossil record.

"The appearance of the first modern primates in North America cooccurred with the appearance of other modern mammals such as horses, and it's all associated with a major global warming event," said co-author Stephen Chester, a Yale University doctoral student and research associate at UF. "It in part set the stage for what we see today in terms of modern mammalian biodiversity."

Less than 6 inches long, *T. brandti* was omnivorous, Bloch said. While archaic primates mostly had claws, some of the characteristics of modern primates include forward-facing eyes, an enlarged brain and nails on all digits.

"They are the smallest true nails known on record, whether living or



fossil," said first author Ken Rose, a professor in the Center for Functional Anatomy & Evolution at Johns Hopkins University School of Medicine. "That certainly doesn't suggest <u>nails</u> developed with larger bodies."

Based on the age of the fossils and analyses of Teilhardina species from other parts of the world, researchers were also able to analyze the hypothesis that mammals migrated from Asia into North America. Instead, they likely passed from Asia, through Europe and into North America on high-latitude land connections.

"This research really suggests that we are looking at something extremely close [to the species found in Europe] and that's of great interest in itself," Rose said. "We can show these species were extremely close morphologically in time and found in Europe and Wyoming."

During the Paleocene-Eocene Thermal Maximum, average temperatures were about 15 degrees Fahrenheit higher than today, and the large variety of mammals found in the fossil record from that time remains a mystery to scientists.

"The finding of this animal and the concentrated effort of this period of time might be one of those things where the closer you look, the less you know," said Gregg Gunnell, director of the Division of Fossil Primates at the Duke Lemur Center. "But any time we have the opportunity to add more morphological information to analyze the relationships of animals to answer these biogeographic questions, we can hopefully get closer and closer to an understanding of what led to this big radiation (diversification) of <u>primates</u> in the first place."

Provided by University of Florida



Citation: Researchers discover oldest evidence of nails in modern primates (2011, August 15) retrieved 1 May 2024 from

https://phys.org/news/2011-08-oldest-evidence-modern-primates.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.