

Shells slim down with CO₂

August 9 2011



Marine algae that turn carbon dissolved in seawater into shell will produce thinner and thinner shells as carbon dioxide levels increase.

The algae, called coccolithophores, have floated in our oceans for over 200 million years Hoovering up carbon and turning it into coccoliths - overlapping plates of [calcium carbonate](#).

Predicting how these algae, an important part of the carbon cycle, will react to rising [CO₂ levels](#) has always been a puzzle. Now a team including Ros Rickaby from Oxford University's Department of Earth Sciences, has found strong evidence that as CO₂ concentration in seawater increases so calcification decreases and coccolith mass declines.

The findings, reported in a recent *Nature* paper, suggest that entire communities of marine organisms, such as coral, are threatened by rising CO₂ and ocean acidification.

The new evidence comes from studies of half a million coccoliths from hundreds of seawater samples and ancient marine sediments cores taken from all over the world.

The research shows much greater variations in coccolith mass than previous lab-based studies, as, in the ocean, rising CO₂ causes populations of algae to favour smaller, lightly calcified species over heavily calcified ones.

Further work is now needed to understand how the algae will respond to the changing marine environment and what impact a rise in thinner-shelled species will have on our oceans and the planet.

More information: *Nature* 476, 80–83 (04 August 2011)
[doi:10.1038/nature10295](https://doi.org/10.1038/nature10295)

Provided by Oxford University

Citation: Shells slim down with CO₂ (2011, August 9) retrieved 30 April 2024 from
<https://phys.org/news/2011-08-shells-slim-co2.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
