


Earthquakes generate big heat in super-small areas: study

October 13 2011



Computer-simulated topography shows high points -- asperities (in red) -- on the rock surface. When in contact with asperities on the adjacent surface, these asperities may undergo intense flash heating in an earthquake. Credit: Mark Robbins and Sangil Hyun, Johns Hopkins University

Most earthquakes that are seen, heard, and felt around the world are caused by fast slip on faults. While the earthquake rupture itself can travel on a fault as fast as the speed of sound or better, the fault surfaces behind the rupture are sliding against each other at about a meter per second.


But the mechanics that underlie fast slip during earthquakes have eluded scientists, because it's difficult to replicate those conditions in the laboratory. "We still largely don't understand what is going at earthquake slip speeds," said David Goldsby, a geophysicist at Brown, "because it's difficult to do experiments at these speeds."

Now, in experiments mimicking earthquake slip rates, Goldsby and Brown geophysicist Terry Tullis show that [fault](#) surfaces in earthquake zones come into contact only at microscopic points between scattered bumps, called asperities, on the fault. These tiny contacts support all the force across the fault. The experiments show that when two fault surfaces slide against other at fast slip rates, the asperities may reach temperatures in excess of 2,700 degrees [Fahrenheit](#), lowering their friction, the scientists write in a paper published in *Science*. The localized, intense heating can occur even while the temperature of the rest of the fault remains largely unaffected, a phenomenon known as flash heating.

"This study could explain a lot of the questions about the mechanics of the [San Andreas Fault](#) and other earthquakes," said Tullis, professor emeritus of [geological sciences](#), who has studied earthquakes for more than three decades.

The experiments simulated earthquake speeds of close to half a meter per second. The rock surfaces touched only at the asperities, each with a surface area of less than 10 microns — a tiny fraction of the total [surface area](#). When the surfaces move against each other at high slip rates, the experiments revealed, heat is generated so quickly at the contacts that temperatures can spike enough to melt most rock types associated with earthquakes. Yet the intense heat is confined to the contact flashpoints; the temperature of the surrounding rock remained largely unaffected by these microscopic hot spots, maintaining a "room temperature" of around 77 degrees Fahrenheit, the researchers write.

"You're dumping in heat extremely quickly into the contacts at high slip rates, and there's simply no time for the heat to get away, which causes the dramatic spike in temperature and decrease in friction," Goldsby said.



Geophysicists Terry Tullis, left, and David Goldsby have shown that rock surfaces sliding past each other in an earthquake can create intense heat but only at the pinpoint places where their surfaces actually touch. Credit: Mike Cohea, Brown University

"The friction stays low so long as the slip rate remains fast," said Goldsby, associate professor of geological sciences (research). "As slip slows, the friction immediately increases. It doesn't take a long time for the fault to restrengthen after you weaken it. The reason is the population of asperities is short-lived and continually being renewed, and therefore at any given slip rate, the asperities have a temperature and therefore friction appropriate for that slip rate. As the slip rate decreases, there is more time for heat to diffuse away from the asperities, and they therefore have lower temperature and higher friction."

Flash heating and other weakening processes that lead to low friction during earthquakes may explain the lack of significant measured heat flows along some active faults like the San Andreas Fault, which might be expected if friction was high on faults during earthquakes. Flash heating in particular may also explain how faults rupture as "slip pulses," wrinkle-like zones of [slip](#) on faults, which would also decrease the

amount of heat generated.

If that is the case, then many earthquakes have been misunderstood as high-friction events. "It's a new view with low dynamic [friction](#). How can it be compatible with what we know?" asked Tullis, who chairs the National [Earthquake](#) Prediction Evaluation Council, an advisory body for the U.S. Geological Survey.

"Flash heating may explain it," Goldsby replied.

Provided by Brown University

Citation: Earthquakes generate big heat in super-small areas: study (2011, October 13) retrieved 26 April 2024 from <https://phys.org/news/2011-10-earthquakes-big-super-small-areas.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--