

Mexican astronomers suggest Bonilla sighting might have been a very close comet breaking up

October 19 2011, by Bob Yirka



Schwassmann-Wachmann3 Comet: Fragment B photographed during its passing in 2006. We can observe up to 73 fragments. This photo is from APOD (Astronomy Picture of the Day) (Internet 3). Credit: NASA, ESA, H. Weaver (JHU/APL), M. Mutchler and Z. Levay (STScl)

(PhysOrg.com) -- Mexican astronomers Hector Javier Durand Manterola, Maria de la Paz Ramos Lara, and Guadalupe Cordero working out of National Autonomous University of Mexico in Mexico City, have uploaded a paper onto the prepress server *arXiv*, suggesting that objects Mexican astronomer José Bonilla recorded in 1883 passing



in front of the sun and surrounded by a hazy mist, were in fact fragments of a comet as it came apart very near the earth.

Binilla wrote down his observations at the time and took some photographs of what he saw, but never really tried to explain what the objects were. Three years later the French astronomy journal *L'Astronomie* published the photos but deemed them likely high flying birds or bugs on the camera lens.

Now Manterola and his colleagues suggest that there is sufficient evidence to at least consider the possibility that the objects Bonilla saw were comet fragments. First of all they say, Bonilla wrote down everything about the sighting; where in the sky the objects appeared, how many fragments he counted (447) over the two days that he could see them, the type of camera he used and the photography techniques, etc. And perhaps most importantly, the time it took for a single fragment to cross the sun, because it's that number that allowed the modern researchers to calculate their distance from Earth, which they suggest would have been somewhere near just 8000 kilometers. Also, because of Bonilla's precision in recording the events, it seems doubtful that he would have overlooked bugs on his lens, or the possibility of the objects being birds, etc.

The unfortunate thing in all this is that no one else reported seeing the objects that Bonilla saw, and there was no associated meteor shower as would be expected if a comet fell apart so close to Earth. Manterola and his team say that this could be because of the parallax effect, which happens when an object is so close, only those closest to it can see it. They also note that because of the apparent position of the objects, they would have been observable only from areas in roughly the same latitude, which would have included parts of Africa and Northern India and south-east Asia; places where no one was looking at the sun through a telescope.



Others clearly are skeptical, for obvious reasons. But that may or may not matter because if the objects were in fact <u>comet</u> fragments, it appears we came pretty close to being wiped out. The researchers suggest that many of the fragments would have been on the order of the size of the one that struck Siberia back in 1908, destroying everything for miles around. If we'd been hit with hundreds or thousands of such fragments at the same time, it might have knocked the life out of our planet, literally.

More information: Interpretation of the observations made in 1883 in Zacatecas (Mexico): A fragmented Comet that nearly hits the Earth, arXiv:1110.2798v1 [astro-ph.EP] <u>arxiv.org/abs/1110.2798</u>

Abstract

In 1883, on the 12th and 13th of August, Mexican astronomer Jose A. y Bonilla observed several objects passing in front of the solar disk. In 1886 in the L'Astronomie magazine, he reported his observations without providing a hypothesis explaining the registered phenomena. Our objective in this work is to interpret, with current knowledge, what he observed in Zacatecas. Our working hypothesis is that what Bonilla observed in 1883 was a highly fragmented comet, in an approach almost flush to the Earth's surface. The fragmentation of the comet's nucleus is a phenomenon known since the XIX century. Using the results reported by Bonilla, we can estimate the distance at which the objects approach to the Earth's surface, their size, their mass and total mass of the comet before fragmentation. According to our calculations, the distance at which the objects passed over the Earth's surface, was between 538 km and 8,062 km, the width of the objects was between 46 m and 795 m and its length between 68 m and 1,022 m, the object's mass was between 5.58e8 kg and 2.5e12 kg. Finally, the mass of the original comet, before fragmentation, was between 1.83e12 and 8.19e15 kg, i.e., between 2e-3 and 8.19 times the mass of Halley Comet.



via TechnologyReview, Discover

© 2011 PhysOrg.com

Citation: Mexican astronomers suggest Bonilla sighting might have been a very close comet breaking up (2011, October 19) retrieved 17 July 2024 from <u>https://phys.org/news/2011-10-mexican-astronomers-bonilla-sighting-comet.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.