

Simple nerve cells regulate swimming depth of marine plankton

October 18 2011



Light microscope image of the larva of the marine annelid *Platynereis*. The larvae swim freely in the sea, moved by activity of their thousands of tiny hair-like structures, which form a band along the larval body (ciliary band), beating coordinately. © Markus Conzelmann, MPI for Developmental Biology


As planktonic organisms the larvae of the marine annelid *Platynereis* swim freely in the open water. They move by activity of their cilia, thousands of tiny hair-like structures forming a band along the larval body and beating coordinately. With changing environmental conditions the larvae swim upward and downward to their appropriate water depth. Scientists of the Max Planck Institute for Developmental Biology in Tübingen, Germany have now identified some signalling substances in the larval nervous system regulating swimming depth of the larvae. These substances influence the ciliary beating and thus hold the larvae in the preferred water depth. The scientists discovered a very simple circuitry of nerve cells underlying this regulation, reflecting an early evolutionary state of the nervous system.

The locomotory system of many animals is muscle based. However, small marine animals often move by cilia. This type of locomotion is more ancient in evolution than muscle-based locomotion and very common in marine plankton. Besides the annelid larvae, the larvae of many marine invertebrates are part of this plankton, for example larvae of snails, sea shells and starfish.

“Not much is known about how the nervous systems of the marine plankton regulate ciliary beating, since the locomotion of intensely explored model organisms like the fruit fly is based on muscles,” says Gáspár Jékely. Together with his team at the Max Planck Institute for [Developmental Biology](#) and in cooperation with Thomas A. Münch at the Centre for Integrative Neuroscience in Tübingen, he has examined in detail the nervous system of marine annelid larvae of *Platynereis dumerilii*.

The ciliary band of *Platynereis* larvae serves as a swimming motor in the seawater: When cilia beat fast and continuously, larvae swim upward, and when cilia cease beating, the larvae sink. These larvae sense different [environmental conditions](#), e.g. they react to changes in

temperature, light and food supply, and alter their movement in the water column accordingly.



Researchers discovered various neuropeptides in the nerve cells of *Platynereis* (white). They are highlighted in different colours in this image. © Albina Asadulina and Markus Conzelmann, MPI for Developmental Biology

In order to gain insight into the regulation of this behaviour, the Tübingen [scientists](#) analysed the genes of *Platynereis*. They discovered

several neuronal signalling substances, so-called neuropeptides in their *Platynereis* gene databases. Moreover, the scientists found that these neuropeptides are produced in single sensory [nerve cells](#) of the larva and are released directly at the ciliary band. The scientists concluded that these nerve cells send the sensory information directly on to the cilia. Some of these neuropeptides influence over cilia beating frequency, others act on the frequency of cilia holdups as well. By means of the neuropeptides, the scientists could control the up and down movement of freely swimming [larvae](#) and change their swimming depth in the water column deliberately.

“We have discovered that the responsible nervous circuitries are built in an unusually simple way. The sensory nerve cells have motor function at the same time, that is, they send the motion impulse directly to the ciliary band,” says Markus Conzelmann from the Max Planck Institute for Developmental Biology, first author of the study. Such simple circuitries are not known from the regulation of muscle-based locomotion. “We were astonished to find not only one neuropeptide as part of such a simple circuitry, but eleven different ones.”

According to the scientists this discovery gives insights into the form and function of nerve systems in an early stage of evolution. Moreover, the results could be interesting for other fields of marine biology: “We now have a suitable model to further explore the regulation of swimming depth in [marine plankton](#). Since the swimming behaviour of plankton is crucial for the survival and prevalence of thousands of marine animal species, our research results could be relevant for marine ecology,” explains Gáspár Jékely. In his future research he wants to reveal how single nerve cells process the different sensory information from water pressure, temperature or salinity.

More information: Markus Conzelmann, Sarah-Lena Offenburger, Albina Asadulina, Timea Keller, Thomas A. Münch and Gáspár Jékely,

Neuropeptides regulate swimming depth of Platynereis larvae. *PNAS*,
[doi: 10.1073/pnas.1109085108](https://doi.org/10.1073/pnas.1109085108)

Provided by Max-Planck-Gesellschaft

Citation: Simple nerve cells regulate swimming depth of marine plankton (2011, October 18)
retrieved 23 May 2024 from <https://phys.org/news/2011-10-simple-nerve-cells-depth-marine.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.