

Subaru's 3-D view of Stephan's Quintet

October 27 2011



Figure 1: Composite tricolor images of Stephan's Quintet using H α filters with a recession velocity of 0 (left image) and a recession velocity of 4,200 miles per second (right image).

Subaru Telescope has added another dimension of information about one of the most studied of all compact galaxy groups—Stephan's Quintet. Located within the borders of the constellation Pegasus, Stephan's Quintet consists of a visual grouping of five galaxies, four of which form an actual compact group of galaxies; one additional galaxy appears in images of the group but is much closer than the others. Refinements in observations of the quintet are revealing more about its members.

A comparison of images (the left and right images in Figure 1) compiled by using a suite of specialized filters with Subaru's Prime Focus Camera (Suprime-Cam) have shown different types of star-formation activity between the closer galaxy NGC7320 and the more distant galaxies in Stephan's Quintet. They show the quintet in 3-D.

These new images are the product of Suprime-Cam's ability to capture images of objects in a wide field of view and to use specialized filters to focus observations according to particular research objectives. To learn about the star-forming regions in Stephan's Quintet and their structures, observers used special narrowband filters for H α emissions, which let in a very specific wavelength of light to indicate distinctive hydrogen emissions during active star formation. They used two H α filters, each with a different recession velocity, i.e. the speed at which the object is moving away from the observer. They used one H α filter with a recession velocity of 0, which means that the speed at which the object is moving away from the observer is 0 and that it is not far distant. They used another H α filter with a greater recession velocity of 4200 miles (6,700 km) per second, an indicator of distant objects. In addition to the red color attributed to the H α emission, blue and green colors assigned to the images from the blue and red filters captured light so that the composite tricolor images aligned with human color perception in red, green, and blue.




Figure 2: A diagram of the member galaxies of Stephan's Quintet. NGC7320 is a closer galaxy and has a recession velocity of 0. The remaining four are a group of more distant galaxies 300 million light years away. The researchers believe

that the merging of NGC7318A/B and NGC7319's crashing into them are responsible for the active star formation regions in the H α emitting region around NGC7318A/B.

Processing of the filtered images resulted in the two different views of Stephan's Quintet shown in Figure 1. The image on the left shows the galaxies when the observers used the H α filter with a recession velocity of 0 while the one on the right shows them when they used the H α filter with a recession velocity of 4,200 miles per second. The left image shows H α emissions that indicate an active star-forming region in the spiral arms of NGC7320 in the lower left quadrant but not in the other galaxies. The right image contrasts with the left and shows a region of H α emissions in the upper three galaxies but none from NGC7320. Two (NGC7318A and NGC7318B) of the four galaxies are shedding gas because of a collision while a third (NGC7319) is crashing in, creating shock waves that trigger vigorous star formation. Figure 2 depicts the relationship of the galaxies. Gas stripped from these three galaxies during galactic collisions is ionized by two mechanisms: shock waves and strong ultraviolet light emanating from the newborn stars. This ionized gas emits bright light, which the H α filter reveals. Thus the researchers believe that NGC7319 as well as NGC7318A/B are driving the star-forming regions in the H α emitting region around NGC7318A/B.

In addition to star-forming activity, the images indicate the distances of the galaxies. Different recession velocities help observers spot cases where objects located at different distances appear in proximity in the same image. The contrasting images show that NGC7320 is closer than the other galaxies, which show active [star formation](#) at a significantly higher recession velocity (4,200 miles per second) than NGC7320 (0). NGC7320 is about 50 million light years away while the other four

galaxies are about 300 million light years away. This explains the intriguing arrangement of the [galaxies](#) in Stephan's Quintet.

Provided by Subaru Telescope

Citation: Subaru's 3-D view of Stephan's Quintet (2011, October 27) retrieved 19 April 2024 from <https://phys.org/news/2011-10-subaru-d-view-stephan-quintet.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.