

# The preferences of uranium: Radionuclide's adsorption in Hanford Site sediments varies based on grain size

October 25 2011

---



More than 200,000 kg of uranium have been released to the vadose zone at the Hanford Site. Plumes of uranium exist in the groundwater and vadose zone.

(PhysOrg.com) -- Uranium prefers petite particles. The radionuclide attaches quickly and abundantly to smaller subsurface grains, according to scientists at Pacific Northwest National Laboratory. The team found that gravel and other large bits adsorbed less uranium(IV) than smaller grains. The larger particles also adsorbed uranium more slowly than the smaller ones. Using this information, the team wrote a series of mathematical formulas to predict uranium adsorption and desorption

affinity and kinetics in sediments containing different grain sizes. The predictions were successfully tested using sediment from the Hanford Site.

Uranium movement at the Hanford Site is an ongoing concern. Uranium and other radioactive holdovers from weapons production reside under the Site, in southeastern Washington State. These [radionuclides](#) are spreading out from their sources and migrating toward the [Columbia River](#), a major waterway. To keep the uranium from the river, scientists need to know how uranium moves, or doesn't, through the complex subsurface. This study demonstrates a way for getting such information for wide areas. With these and other results, PNNL scientists are shaping how uranium's behavior is considered by scientists, cleanup experts, and regulators.

Calculating exactly how much uranium is present and how fast uranium migrates requires scientists to scale measurements from the lab to the field. Essentially, they must take samples measured across millimeters to centimeters and extrapolate them to systems that extend for meters or kilometers.

"Upscaling is always a challenge," said Dr. Chongxuan Liu, a PNNL [geochemist](#) who worked on the study.

The scientists on this study began by conducting [wet chemistry](#) experiments to determine adsorption characteristics of four grain sizes. The grains were coarse sand (1-2 mm), medium sand (0.2-1 mm), fine sand (0.053-0.2 mm), and clay/silt fraction (

Citation: The preferences of uranium: Radionuclide's adsorption in Hanford Site sediments varies based on grain size (2011, October 25) retrieved 20 April 2024 from <https://phys.org/news/2011-10-uranium-radionuclide-adsorption-hanford-site.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.