

Study describes how space flight impacts astronauts' eyes and vision

November 3 2011

A new study sponsored by NASA finds that space flights lasting six months or more can cause a spectrum of changes in astronauts' visual systems. Some problems, including blurry vision, appear to persist long after astronauts' return to Earth. The results are affecting plans for longduration manned space voyages, such as a trip to Mars. The study team included ophthalmologists Thomas H. Mader, MD, of Alaska Native Medical Center and Andrew G. Lee, MD, of The Methodist Hospital, Houston, Texas. Their report is published in October's *Ophthalmology*, the journal of the American Academy of Ophthalmology.

The researchers studied seven <u>astronauts</u>, all of whom were about age 50 and had spent at least six continuous months in space. All reported that their <u>vision</u> became blurry, to varying degrees, while on the space station. Vision changes usually began around six weeks into the mission and persisted in some astronauts for months after their return to Earth. Drs. Mader and Lee agree that the eye abnormalities appear to be unrelated to launch or re-entry, since they occurred only in astronauts who spent extended time in microgravity.

In-depth examination of the seven astronauts revealed several abnormalities. All of the subjects had one or more of the following changes in the tissues, fluids, nerves and other structures in the back of the eye:

• Flattening of the back of the eyeball (five subjects);



- Folds in the choroid, the <u>vascular tissue</u> behind the retina, which is the light sensitive area in the back of the eye (five subjects); and
- Excess fluid around and presumed swelling of the optic nerve (five subjects).

Such abnormalities could potentially be caused by increased intracranial pressure-that is, pressure inside the head. However, none of these astronauts experienced symptoms usually associated with intracranial pressure, such as chronic headache, double vision, or ringing in the ears. Researchers believe other factors may be involved, such as abnormal flow of spinal fluid around the optic nerve, changes in blood flow in the choroid, or changes related to chronic low pressure within the eye, which is known as intraocular pressure. They hypothesize that these changes may result from the fluid shifts toward the head that occur when astronauts spend extended time in microgravity.

The visual system changes discovered by the researchers may represent a set of adaptations to microgravity. The degree and type of response appear to vary among astronauts. Researchers hope to discover whether some astronauts are less affected by <u>microgravity</u> and therefore better-suited for extended <u>space flight</u>, such as a three-year round trip to Mars.

In their report, Drs. Mader and Lee also noted a recent NASA survey of 300 astronauts that found that correctible problems with both near and distance vision were reported by about 23 percent of astronauts on brief missions and by 48 percent of those on extended missions. The survey confirmed that for some astronauts, these vision changes continue for months or years after return to Earth. The possibility of near vision problems has been recognized for decades, and special "space anticipation glasses" to improve visual sharpness have been provided on all spacecraft dating back to John Glenn, who had a pair in his space capsule.



"In astronauts over age 40, like non-astronauts of the same age, the eye's lens may have lost some of its ability to change focus," said Dr. Mader. "In the <u>space</u> program's early days most astronauts were younger, military test-pilots who had excellent vision. Today's astronauts tend to be in their 40s or older. This may be one reason we've seen an uptick in vision problems. Also, we suspect many of the younger astronauts were more likely to 'tough out' any problems they experienced, rather than reporting them."

As part of ongoing research all astronauts now receive comprehensive eye exams and vision testing. Diagnostic tests include pre- and postflight magnetic resonance imaging, optical coherence tomography, which magnifies cross-section views of parts of the eye, and fundus photography, which records images of the retina and back of the eye. Intraocular pressure measurement and ultrasound imaging take place in flight, as well as pre- and post-mission.

Provided by American Academy of Ophthalmology

Citation: Study describes how space flight impacts astronauts' eyes and vision (2011, November 3) retrieved 16 May 2024 from <u>https://phys.org/news/2011-11-space-flight-impacts-astronauts-eyes.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.