

Researchers discover novel chemical route to form organic molecules

January 13 2012

An international team of scientists led by University of Hawai'i at Mānoa Professor Ralf I. Kaiser, Alexander M. Mebel of Florida International University, and Alexander Tielens of Leiden Observatory in the Netherlands, discovered a novel chemical route to form polycyclic aromatic hydrocarbons (PAHs) – complex organic molecules such as naphthalene carrying fused benzene rings – in ultra-cold regions of interstellar space. The team announced their findings in the January 3 issue of Proceedings of the National Academy of Sciences. Funding for the study was provided by the U.S. Department of Energy, Basic Energy Sciences.

These findings have crucial implications not only to reduce the emission of PAHs as toxic byproducts from internal combustion engines, but also rationalize the synthetic routes to a key class of [organic molecules](#) in the [interstellar medium](#) associated with the origins of life.

On Earth, PAHs are associated with incomplete combustion processes and can be formed readily at elevated temperatures in combustion engines of cars and in cigarette smoke. Once liberated into the ambient environment, PAHs can be transferred into the lungs by inhalation and are strongly implicated in the degradation of human health, particularly due to their high carcinogenic risk potential. PAHs are also serious water pollutants of marine ecosystems and bioaccumulate in the fatty tissue of living organisms. Together with leafy vegetables, where PAHs deposit easily, they have been further linked to soil contamination, food poisoning, liver lesions, and tumor growth.

Whereas on Earth, PAHs are classified as highly toxic, PAHs have been dubbed as the ‘cradle of life’ in the interstellar medium and are considered as key players in the astrobiological evolution. On the molecular level, functionalized PAHs carrying carbonyl and hydroxyl groups were found in organic extracts from the Murchison meteorite and form membrane-like boundary structures, the first indications of a cell type structure, which are requisite to the origin of life. The compounds that are water soluble form non-soluble vesicles, constituting molecules that possess both polar and non-polar components. The hollow droplets formed by this lipid multilayer are essential for the origin of life process since they provide an environment in which the functionalized PAHs can evolve by isolating and protecting them from the surrounding medium.

[Scientists](#) have been researching the formation of PAHs in combustion flames and in the interstellar medium for decades, but the formation mechanism of even the simplest PAH prototype – the naphthalene molecule ($C_{10}H_8$) as present in earthly mothballs - has remained an open question. Textbook knowledge postulates that classical reaction mechanisms involve complex reactions following hydrogen abstraction and acetylene addition (HACA) sequences with substantial ‘activation energies.’ These processes can only operate at high temperatures of a few 1,000 K as present, for instance, in combustion processes and in the outflows of carbon-rich stars and planetary nebulae. However, in recent years it has become quite clear that interstellar PAHs are rapidly destroyed in the interstellar medium upon photolysis, interstellar shock waves driven by supernova explosions, and energetic cosmic rays. The destruction time scales are much shorter than the timescale for injection of new material into the interstellar medium by carbon-rich Asymptotic Giant Branch (AGB) stars and carbon-rich planetary nebulae as the descendants of AGB stars. Therefore, the ubiquitous presence of PAHs in the interstellar medium implies a crucial, previously unexplained route to a fast chemical growth of PAHs in the cold environment of the interstellar medium at temperatures down to 10 K, where the classical

quinoline. This concept can be also expanded to functionalized PAHs with organic side chains thus bringing researchers closer to solving the decade old puzzle of how complex PAHs and their derivatives can be synthesized in combustion flames and in cold interstellar space.

Provided by University of Hawaii at Manoa

Citation: Researchers discover novel chemical route to form organic molecules (2012, January 13) retrieved 2 May 2024 from <https://phys.org/news/2012-01-chemical-route-molecules.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.